

<b>Table of Contents</b> <b>Section II – County Soil and Site Information Reports</b>
--

	Issue Date	Date of Last Review	Responsible Staff
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
<b>Soils Legends</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Acreage and Proportionate Extent of the Soils</i>			
<b>Soil Descriptions - Nontechnical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
<b>Soil Descriptions - Technical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Map Unit Description Report</i>			
<b>Cropland Interpretations - Technical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Land Capability and Yields Per Acre of Crops</i>			
<i>*Prime Farmland Report</i>			
<i>*Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
<b>Rangeland, Grazed Forestland, Native Pastureland Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
<b>Forestland Interpretations</b>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
<b>Nonagricultural Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
<b>Recreation Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Recreational Interpretations</i>			
<b>Wildlife Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Wildlife Interpretations Report</i>			

<b>Pastureland and Hayland Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Yields Per Acre of Pasture and Hayland</i>			
	<b>Issue Date</b>	<b>Date of Last Review</b>	<b>Responsible Staff</b>
<b>Mined Land Interpretations</b>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Mined Land Interpretations</i>			
<b>Windbreak Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Conservation Tree and Shrub Management Report</i>			
<b>Engineering Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Engineering Index Properties</i>			
<i>*Physical Properties of the Soils</i>			
<i>*Chemical Properties of the Soils</i>			
<i>*Water Features</i>			
<i>*Soil Features</i>			
<i>*Water Management Report</i>			
<b>Waste Disposal Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Sanitary Facilities Report</i>			
<i>*Agricultural Waste Management Report</i>			
<b>Hydric Soil Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>Use and Explanation of Hydric Soil Interpretations</i>			
<i>*Hydric Soils List</i>			
<b>HEL Interpretations</b>	<b>See Report</b>		<b>SOI</b>
<i>Use and Explanation of Highly Erodible Land Interpretations</i>			
<i>*Highly Erodible Lands Report</i>			
1	<i>Highly Erodible Land</i>		
2	<i>Potentially Highly Erodible Land</i>		
3	<i>Nonhighly Erodible Land</i>		

*\*County specific computer generated reports.*

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

Map symbol	Soil name	Acres	Percent
Ab	Albaton Silt Loam, Overwash-----	926	0.2
Ac	Albaton Silty Clay-----	4,424	1.2
Af	Alda Fine Sandy Loam-----	703	0.2
Ag	Alda Very Fine Sandy Loam-----	2,310	0.6
Ca	Carr Fine Sandy Loam-----	1,616	0.4
Cc	Cass Fine Sandy Loam-----	1,805	0.5
Cd	Cass Fine Sandy Loam, Loamy Substratum-----	1,105	0.3
Ce	Cass Very Fine Sandy Loam-----	2,219	0.6
Cg	Colo Silty Clay Loam-----	1,198	0.3
Ck	Colo And Kennebec Soils-----	13,379	3.6
Cm	Cut And Fill Land-----	4,591	1.2
DcE	Dickinson Soils, 11 To 17 Percent Slopes-----	335	*
Ed	Eudora Silt Loam-----	3,948	1.1
Ga	Gibbon Loamy Sand, Overwash-----	1,148	0.3
Gb	Gibbon Silt Loam-----	15,274	4.1
Gc	Gibbon Silty Clay Loam-----	5,123	1.4
Gs	Gibbon-Slickspots Complex-----	266	*
Gu	Gullied Land-----	2,538	0.7
Ha	Haynie Silt Loam-----	2,239	0.6
IdD2	Ida Silt Loam, 7 To 17 Percent Slopes, Eroded-----	2,933	0.8
IdE	Ida Silt Loam, 17 To 30 Percent Slopes-----	1,826	0.5
IdE2	Ida Silt Loam, 17 To 30 Percent Slopes, Eroded-----	1,147	0.3
Im	Inavale Loamy Fine Sand-----	4,142	1.1
In	Inavale Loamy Fine Sand, Hummocky-----	816	0.2
JuB	Judson Silt Loam, 3 To 7 Percent Slopes-----	26,050	6.9
Ke	Kennebec Silt Loam, Occasionally Flooded-----	14,474	3.9
Le	Lex Soils, Noncalcareous Variant-----	2,852	0.8
Ls	Luton Silt Loam, Overwash-----	867	0.2
Lt	Luton Silty Clay Loam-----	1,168	0.3
Lu	Luton Silty Clay-----	616	0.2
MaA	Marshall Silty Clay Loam, 0 To 1 Percent Slopes-----	4,182	1.1
MaB	Marshall Silty Clay Loam, 1 To 3 Percent Slopes-----	1,874	0.5
MaC	Marshall Silty Clay Loam, 3 To 7 Percent Slopes-----	42,264	11.2
MaC2	Marshall Silty Clay Loam, 3 To 7 Percent Slopes, Eroded-----	2,521	0.7
MaD	Marshall Silty Clay Loam, 7 To 11 Percent Slopes-----	19,961	5.3
MeD2	Marshall-Ponca Silty Clay Loams, 7 To 11 Percent Slopes, Eroded-----	47,374	12.6
MfE	Marshall And Ponca Soils, 11 To 17 Percent Slopes-----	3,466	0.9
MfE2	Marshall And Ponca Soils, 11 To 17 Percent Slopes, Eroded-----	21,317	5.7
MoA	Monona Silt Loam, 0 To 1 Percent Slopes-----	2,104	0.6
MoB	Monona Silt Loam, 1 To 3 Percent Slopes-----	751	0.2
MoC	Monona Silt Loam, 3 To 7 Percent Slopes-----	10,674	2.8
MoC2	Monona Silt Loam, 3 To 7 Percent Slopes, Eroded-----	620	0.2
MoD	Monona Silt Loam, 7 To 11 Percent Slopes-----	10,095	2.7
MoD2	Monona Silt Loam, 7 To 11 Percent Slopes, Eroded-----	6,806	1.8
MoE	Monona Silt Loam, 11 To 17 Percent Slopes-----	15,116	4.0
MsE2	Monona And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded-----	11,354	3.0
MsF	Monona And Ida Silt Loams, 17 To 30 Percent Slopes-----	3,733	1.0
MsF2	Monona And Ida Silt Loams, 17 To 30 Percent Slopes, Eroded-----	2,856	0.8
MsG	Monona And Ida Silt Loams, 30 To 60 Percent Slopes-----	2,297	0.6
On	Onawa Silty Clay-----	987	0.3
Pa	Percival Silty Clay-----	947	0.3
Pb	Pits And Dumps-----	2,030	0.5
Pc	Platte Soils-----	2,033	0.5
PdD2	Ponca And Ida Silt Loams, 7 To 11 Percent Slopes, Eroded-----	2,941	0.8
PdE2	Ponca And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded-----	3,319	0.9
Ra	Riverwash-----	1,008	0.3
Rk	Rock Land-----	342	*
Rn	Rough Broken Land, Loess-----	1,132	0.3
Sd	Sandy Alluvial Land-----	5,330	1.4
Sp	Sarpy Fine Sand-----	1,079	0.3
Ss	Silty Alluvial Land-----	1,735	0.5
StE2	Steinauer Clay Loam, 11 To 30 Percent Slopes, Eroded-----	678	0.2
Wb	Wabash Silt Loam-----	918	0.2
Wc	Wabash Silty Clay-----	2,383	0.6
Wm	Wann Fine Sandy Loam-----	4,173	1.1
Wt	Wet Alluvial Land-----	1,705	0.5
zwa	Water < 40 Acres-----	2,769	0.7
zwb	Water > 40 Acres-----	11,240	3.0
	Total-----	378,152	100.6

\* Less than 0.1 percent.



Nontechnical Soil Descriptions  
Douglas and Sarpy Counties, Nebraska

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

---

Ab Albaton Silt Loam, Overwash

Albaton soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of calcareous clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 3w.

Ac Albaton Silty Clay

Albaton soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of calcareous clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Af Alda Fine Sandy Loam

Alda soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Ag Alda Very Fine Sandy Loam

Alda soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Ca Carr Fine Sandy Loam

Carr soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Cc Cass Fine Sandy Loam

Cass soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of sandy and/or loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

Cd Cass Fine Sandy Loam, Loamy Substratum

Cass soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of sandy and/or loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Ce Cass Very Fine Sandy Loam

Cass soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of sandy and/or loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

Cg Colo Silty Clay Loam

Colo soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of fine-silty alluvium. This soil is poorly drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Ck Colo And Kennebec Soils

Colo soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of fine-silty alluvium. This soil is poorly drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Kennebec soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of silty alluvium. This soil is moderately well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 54 inches. This soil is in the Silty Lowland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 2w.

DcE Dickinson Soils, 11 To 17 Percent Slopes

Dickinson soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep bluff on upland. The runoff class is low. The parent material consists of wind reworked glaciofluvial deposits. This soil is somewhat excessively drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Ed Eudora Silt Loam

Eudora soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of coarse-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Silty Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

Ga Gibbon Loamy Sand, Overwash

Gibbon soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

## Gb Gibbon Silt Loam

Gibbon soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

## Gc Gibbon Silty Clay Loam

Gibbon soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

## Gs Gibbon-Slickspots Complex

Gibbon soil makes up 65 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Saltine soil makes up 35 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a very high available water capacity and a high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a moderately saline horizon, it has a horizon that is strongly sodic. This soil is in the Saline Subirrigated - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6s.

## Gu Gullied Land

Ida Variant soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a steep to 0.000000 gully on loess hill on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Thin Loess - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 7e.

## Ha Haynie Silt Loam

Haynie soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of coarse-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

## IdD2 Ida Silt Loam, 7 To 17 Percent Slopes, Eroded

Ida soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

## IdE Ida Silt Loam, 17 To 30 Percent Slopes

Ida soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

IdE2 Ida Silt Loam, 17 To 30 Percent Slopes, Eroded

Ida soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Im Inavale Loamy Fine Sand

Inavale soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

In Inavale Loamy Fine Sand, Hummocky

Inavale soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a moderately sloping flood plain on river valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

JuB Judson Silt Loam, 3 To 7 Percent Slopes

Judson soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately sloping drainage way on upland. The runoff class is low. The parent material consists of fine-silty colluvium. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 2e.

Ke Kennebec Silt Loam, Occasionally Flooded

Kennebec soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of silty alluvium. This soil is moderately well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 54 inches. This soil is in the Silty Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Le Lex Soils, Noncalcareous Variant

Lockton soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a moderate available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Ls Luton Silt Loam, Overwash

Luton soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Lt Luton Silty Clay Loam

Luton soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.



Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

Lu Luton Silty Clay

Luton soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

MaA Marshall Silty Clay Loam, 0 To 1 Percent Slopes

Marshall soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level loess hill on upland. The runoff class is low. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

MaB Marshall Silty Clay Loam, 1 To 3 Percent Slopes

Marshall soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a gently sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

MaC Marshall Silty Clay Loam, 3 To 7 Percent Slopes

Marshall soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a moderately sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 2e.

MaC2 Marshall Silty Clay Loam, 3 To 7 Percent Slopes, Eroded

Marshall soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a moderately sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

MaD Marshall Silty Clay Loam, 7 To 11 Percent Slopes

Marshall soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is high. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

MeD2 Marshall-Ponca Silty Clay Loams, 7 To 11 Percent Slopes, Eroded

Marshall soil makes up 70 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is high. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

Ponca soil makes up 30 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

MfE Marshall And Ponca Soils, 11 To 17 Percent Slopes

Marshall soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

Ponca soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

MfE2 Marshall And Ponca Soils, 11 To 17 Percent Slopes, Eroded

Marshall soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty noncalcareous loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

Ponca soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

MoA Monona Silt Loam, 0 To 1 Percent Slopes

Monona soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level loess hill on upland. The runoff class is negligible. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

MoB Monona Silt Loam, 1 To 3 Percent Slopes

Monona soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a gently sloping loess hill on upland. The runoff class is low. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

MoC Monona Silt Loam, 3 To 7 Percent Slopes

Monona soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately sloping loess hill on upland. The runoff class is low. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 2e.

MoC2 Monona Silt Loam, 3 To 7 Percent Slopes, Eroded

Pohocco soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

MoD Monona Silt Loam, 7 To 11 Percent Slopes

Monona soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

MoD2 Monona Silt Loam, 7 To 11 Percent Slopes, Eroded

Pohocco soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

MoE Monona Silt Loam, 11 To 17 Percent Slopes

Monona soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is medium. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

MsE2 Monona And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded

Pohocco soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

Ida soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep loess hill on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

MsF Monona And Ida Silt Loams, 17 To 30 Percent Slopes

Monona soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Ida soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

MsF2 Monona And Ida Silt Loams, 17 To 30 Percent Slopes, Eroded

Monona soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

Ida soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

MsG Monona And Ida Silt Loams, 30 To 60 Percent Slopes

Monona soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a steep to steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 7e.

Ida soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a steep to very steep loess bluff on upland. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Thin Loess - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 7e.

On Onawa Silty Clay

Onawa soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of calcareous clayey alluvium over loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Pa Percival Silty Clay

Percival soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a low available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Pc Platte Soils

Platte soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

PdD2 Ponca And Ida Silt Loams, 7 To 11 Percent Slopes, Eroded

Ponca soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

Ida soil makes up 50 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping loess hill on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

---

PdE2 Ponca And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded

Ponca soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

Ida soil makes up 50 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a moderately steep loess hill on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 4e.

Ra Riverwash

Riverwash soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified sandy alluvium. This soil is very poorly drained. The slowest permeability is rapid. It has a very low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It is in the nonirrigated land capability classification 8w.

Rn Rough Broken Land, Loess

Ida Variant soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a steep to very steep gully on loess hill on upland. The runoff class is high. The parent material consists of fine-silty loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Thin Loess - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 7e.

Sd Sandy Alluvial Land

Inavale soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy Lowland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6w.

Sp Sarpy Fine Sand

Sarpy soil makes up 100 percent of the map unit. This map unit is in the Iowa and Missouri Deep Loess Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of calcareous sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 2 percent calcium carbonate. This soil is in the Sandy Lowland - Veg. Zone 4 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

Ss Silty Alluvial Land

Nodaway Variant soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is moderately well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 54 inches. This soil is in the Silty Overflow - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6w.

StE2 Steinauer Clay Loam, 11 To 30 Percent Slopes, Eroded

Steinauer soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a strongly sloping to steep hill on upland. The runoff class is very high. The parent material consists of calcareous loamy till. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 20 percent calcium carbonate. This soil is in the Limey Upland - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 6e.

Nontechnical Soil Descriptions--Continued  
Douglas and Sarpy Counties, Nebraska

## Wb Wabash Silt Loam

Wabash soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

## Wc Wabash Silty Clay

Wabash soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is high. The parent material consists of clayey alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a very high shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Clayey Overflow - Veg. Zone 4 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

## Wm Wann Fine Sandy Loam

Wann soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of calcareous loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 30 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 4 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

## Wt Wet Alluvial Land

Barney soil makes up 100 percent of the map unit. This map unit is in the Nebraska and Kansas Loess-Drift Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is very poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land - Veg. Zone 4 range site. It is in the nonirrigated land capability classification 5w.



**Ab—Albaton silt loam, Overwash**

## Map Unit Composition

Albaton: 100 percent

## Component Descriptions

Albaton

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Flood plain on river valley

*Parent material:* Calcareous clayey alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Very slow (About 0.00 in/hr)

*Available water capacity:* Moderate (About 8.3 inches)

*Shrink-swell potential:* Very high (About 17.0 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 0 to 18 inches

*Runoff class:* High

*Ecological site:* Clayey Overflow - Veg. Zone 4

*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 18 inches; silt loam

H2—18 to 60 inches; silty clay

*Shrink-swell potential:* Very high (About 17.0 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 0 to 18 inches

*Runoff class:* High

*Ecological site:* Clayey Overflow - Veg. Zone 4

*Land capability (irrigated):* 3w

*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 6 inches; silty clay

H2—6 to 60 inches; silty clay

**Af—Alda fine sandy loam**

## Map Unit Composition

Alda: 100 percent

## Component Descriptions

Alda

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Low (About 4.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 3w

*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 13 inches; fine sandy loam

H2—13 to 20 inches; fine sandy loam

H3—20 to 60 inches; coarse sand

**Ac—Albaton silty clay**

## Map Unit Composition

Albaton: 100 percent

## Component Descriptions

Albaton

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Flood plain on river valley

*Parent material:* Calcareous clayey alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Very slow (About 0.00 in/hr)

*Available water capacity:* Moderate (About 6.0 inches)

**Ag—Alda very fine sandy loam**

## Map Unit Composition

Alda: 100 percent



**Component Descriptions**

Alda

MLRA: 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley*Parent material:* Loamy alluvium over sandy and gravelly alluvium*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Low (About 5.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Negligible*Ecological site:* Subirrigated - Veg. Zone 4*Land capability (irrigated):* 3w*Land capability (nonirrigated):* 3w**Typical Profile:**

H1—0 to 13 inches; very fine sandy loam

H2—13 to 20 inches; fine sandy loam

H3—20 to 60 inches; coarse sand

**Ca—Carr fine sandy loam****Map Unit Composition**

Carr: 100 percent

**Component Descriptions**

Carr

MLRA: 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Flood plain on river valley*Parent material:* Stratified calcareous alluvium*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 9.6 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 4*Land capability (irrigated):* 2w*Land capability (nonirrigated):* 2w**Typical Profile:**

H1—0 to 15 inches; fine sandy loam

H2—15 to 60 inches; stratified fine sandy loam to loam

**Cc—Cass fine sandy loam****Map Unit Composition**

Cass: 100 percent

**Component Descriptions**

Cass

MLRA: 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley*Parent material:* Sandy and/or loamy alluvium*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Moderate (About 7.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Rare*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 4*Land capability (irrigated):* 2e*Land capability (nonirrigated):* 2e**Typical Profile:**

H1—0 to 16 inches; fine sandy loam

H2—16 to 39 inches; fine sandy loam

H3—39 to 60 inches; coarse sand

**Cd—Cass fine sandy loam, loamy Substratum****Map Unit Composition**

Cass: 100 percent

**Component Descriptions**

Cass

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Sandy and/or loamy alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 8.8 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* About 30 to 42 inches

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland - Veg. Zone 4

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 20 inches; fine sandy loam

H2—20 to 30 inches; fine sandy loam

H3—30 to 48 inches; silty clay loam

H4—48 to 60 inches; coarse sand

## **Ce—Cass very fine sandy loam**

### Map Unit Composition

Cass: 100 percent

### Component Descriptions

Cass

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Sandy and/or loamy alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland - Veg. Zone 4

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 1

*Typical Profile:*

H1—0 to 16 inches; very fine sandy loam

H2—16 to 39 inches; fine sandy loam

H3—39 to 60 inches; coarse sand

## **Cg—Colo silty clay loam**

### Map Unit Composition

Colo: 100 percent

### Component Descriptions

Colo

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Fine-silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.9 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 0 to 18 inches

*Runoff class:* Low

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 2w

*Typical Profile:*

H1—0 to 23 inches; silty clay loam

H2—23 to 40 inches; silty clay loam

H3—40 to 60 inches; silty clay loam

## **Ck—Colo And Kennebec Soils**

### Map Unit Composition

Colo: 50 percent

Kennebec: 50 percent

### Component Descriptions

Colo

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley  
*Parent material:* Fine-silty alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.9 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* Low  
*Ecological site:* Subirrigated - Veg. Zone 4  
*Land capability (irrigated):* 2w  
*Land capability (nonirrigated):* 2w

*Typical Profile:*

H1—0 to 23 inches; silty clay loam  
H2—23 to 40 inches; silty clay loam  
H3—40 to 60 inches; silty clay loam

**Kennebec**

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley  
*Parent material:* Silty alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Moderately well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 11.9 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 36 to 72 inches  
*Runoff class:* Negligible  
*Ecological site:* Silty Lowland - Veg. Zone 4  
*Land capability (nonirrigated):* 2w

*Typical Profile:*

H1—0 to 17 inches; silt loam  
H2—17 to 60 inches; silt loam

## **Cm—Cut And Fill Land**

**Map Unit Composition**

Urban land: 100 percent

**Component Descriptions**

Urban land

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Hill on upland, drainageway on upland

*Slope:* 0 to 30 percent

*Available water capacity:* Very low (About 0.0 inches)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very high

## **DcE—Dickinson Soils, 11 to 17 percent slopes**

**Map Unit Composition**

Dickinson: 100 percent

**Component Descriptions**

Dickinson

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Bluff on upland

*Parent material:* Wind reworked glaciofluvial deposits

*Slope:* 11 to 17 percent

*Drainage class:* Somewhat excessively drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 10 inches; fine sandy loam  
H2—10 to 16 inches; fine sandy loam  
H3—16 to 30 inches; fine sandy loam  
H4—30 to 60 inches; loamy sand

## **Ed—Eudora silt loam**

**Map Unit Composition**

Eudora: 100 percent

**Component Descriptions**

Eudora

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Coarse-silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Silty Overflow - Veg. Zone 4

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 1

**Typical Profile:**

H1—0 to 13 inches; silt loam

H2—13 to 60 inches; silt loam

## **Ga—Gibbon loamy sand, Overwash**

**Map Unit Composition**

Gibbon: 100 percent

**Component Descriptions**

Gibbon

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Stratified calcareous silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 9.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 3w

*Land capability (nonirrigated):* 3w

**Typical Profile:**

H1—0 to 17 inches; loamy sand

H2—17 to 60 inches; loam

## **Gb—Gibbon silt loam**

**Map Unit Composition**

Gibbon: 100 percent

**Component Descriptions**

Gibbon

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Stratified calcareous silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderate (About 0.57 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 2w

**Typical Profile:**

H1—0 to 16 inches; silt loam

H2—16 to 40 inches; silt loam

H3—40 to 60 inches; stratified very fine sandy loam to silt loam

## **Minor Components Ponded Soils**

## **Gc—Gibbon silty clay loam**

**Map Unit Composition**

Gibbon: 100 percent

**Component Descriptions**

Gibbon

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Stratified calcareous silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Low

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 2w

**Typical Profile:**

H1—0 to 16 inches; silty clay loam

H2—16 to 40 inches; silty clay loam

H3—40 to 60 inches; stratified very fine sandy loam to silty clay loam

**Minor Components**

**Wt At 0-1 Foot**

**Gs—Gibbon-Slickspots complex****Map Unit Composition**

Gibbon: 65 percent

Saltine: 35 percent

**Component Descriptions**

Gibbon

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Stratified calcareous silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 2w

**Typical Profile:**

H1—0 to 20 inches; silt loam

H2—20 to 40 inches; silty clay loam

H3—40 to 60 inches; stratified very fine sandy loam to silty clay loam

Saltine

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.9 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Low

*Ecological site:* Saline Subirrigated - Veg. Zone 4

*Land capability (nonirrigated):* 6s

**Typical Profile:**

H1—0 to 6 inches; silt loam

H2—6 to 15 inches; silty clay loam

H3—15 to 20 inches; silty clay loam

H4—20 to 60 inches; silty clay loam

**Minor Components**

**Wt At 0-1 Foot**

**Gu—Gullied land****Map Unit Composition**

Ida Variant: 100 percent

## Component Descriptions

Ida Variant

MLRA: 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Gully on loess hill on upland*Parent material:* Fine-silty calcareous loess*Slope:* 30 to 99 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 11.3 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* High*Ecological site:* Thin Loess - Veg. Zone 4*Land capability (nonirrigated):* 7e*Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 60 inches; silt loam

**Ha—Haynie silt loam**

## Map Unit Composition

Haynie: 100 percent

## Component Descriptions

Haynie

MLRA: 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Flood plain on river valley*Parent material:* Coarse-silty alluvium*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* Very high (About 12.4 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Silty Lowland - Veg. Zone 4*Land capability (irrigated):* 2w*Land capability (nonirrigated):* 2w*Typical Profile:*

H1—0 to 9 inches; silt loam

H2—9 to 60 inches; silt loam

**IdD2—Ida silt loam, 7 to 17 percent slopes, Eroded**

## Map Unit Composition

Ida: 100 percent

## Component Descriptions

Ida

MLRA: 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland*Parent material:* Fine-silty calcareous loess*Slope:* 7 to 17 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* Very high (About 12.4 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Medium*Ecological site:* Limy Upland - Veg. Zone 4*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

**IdE—Ida silt loam, 17 to 30 percent slopes**

## Map Unit Composition

Ida: 100 percent

## Component Descriptions

Ida

MLRA: 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland*Parent material:* Fine-silty calcareous loess*Slope:* 17 to 30 percent

*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.4 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* High  
*Ecological site:* Limy Upland - Veg. Zone 4  
*Land capability (nonirrigated):* 6e

*Typical Profile:*  
 H1—0 to 5 inches; silt loam  
 H2—5 to 60 inches; silt loam

## **IdE2—Ida silt loam, 17 to 30 percent slopes, Eroded**

Map Unit Composition

Ida: 100 percent

Component Descriptions

Ida

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 17 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Limy Upland - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*  
 H1—0 to 5 inches; silt loam  
 H2—5 to 60 inches; silt loam

## **Im—Inavale loamy fine sand**

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Sandy alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland - Veg. Zone 4

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 11 inches; loamy fine sand

H2—11 to 28 inches; loamy fine sand

H3—28 to 60 inches; fine sand

## **Minor Components Ponded Soils**

## **In—Inavale loamy fine sand, Hummocky**

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Sandy alluvium

*Slope:* 3 to 7 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 8 inches; loamy fine sand

H2—8 to 25 inches; loamy fine sand

H3—25 to 60 inches; fine sand

## **JuB—Judson silt loam, 3 to 7 percent slopes**

Map Unit Composition

Judson: 100 percent

Component Descriptions

Judson

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Drainageway on upland

*Parent material:* Fine-silty colluvium

*Slope:* 3 to 7 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.40 in/hr)

*Available water capacity:* High (About 12.0 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 26 inches; silt loam

H2—26 to 37 inches; silty clay loam

H3—37 to 60 inches; silty clay loam

### **Minor Components**

**Wt At 0-1 Foot**

## **Ke—Kennebec silt loam, Occasionally Flooded**

Map Unit Composition

Kennebec: 100 percent

Component Descriptions

Kennebec

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Moderately well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.3 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 36 to 72 inches

*Runoff class:* Negligible

*Ecological site:* Silty Overflow - Veg. Zone 4

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 2w

*Typical Profile:*

H1—0 to 26 inches; silt loam

H2—26 to 60 inches; silt loam

### **Minor Components**

**Ponded Soils**

## **Le—Lex Soils, Noncalcareous Variant**

Map Unit Composition

Lockton: 100 percent

Component Descriptions

Lockton

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Loamy alluvium

*Slope:* 0 to 1 percent



*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Moderate (About 6.7 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Subirrigated - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 21 inches; silt loam  
 H2—21 to 28 inches; sandy loam  
 H3—28 to 60 inches; gravelly sand

**Minor Components**  
**Wt At 0-1 Foot**

**Ls—Luton silt loam, Overwash**

Map Unit Composition

Luton: 100 percent

Component Descriptions

Luton  
 MLRA: 107 - Iowa and Missouri Deep Loess Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Clayey alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Very slow (About 0.00 in/hr)  
*Available water capacity:* Moderate (About 7.5 inches)  
*Shrink-swell potential:* Very high (About 17.0 LEP)  
*Flooding hazard:* Rare  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* High  
*Ecological site:* Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 11 inches; silt loam

H2—11 to 27 inches; silty clay  
 H3—27 to 60 inches; silty clay

**Lt—Luton silty clay loam**

Map Unit Composition

Luton: 100 percent

Component Descriptions

Luton  
 MLRA: 107 - Iowa and Missouri Deep Loess Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Clayey alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Very slow (About 0.00 in/hr)  
*Available water capacity:* Moderate (About 7.1 inches)  
*Shrink-swell potential:* Very high (About 17.0 LEP)  
*Flooding hazard:* Rare  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* High  
*Ecological site:* Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 13 inches; silty clay loam  
 H2—13 to 27 inches; silty clay  
 H3—27 to 60 inches; silty clay

**Lu—Luton silty clay**

Map Unit Composition

Luton: 100 percent

Component Descriptions

Luton  
 MLRA: 107 - Iowa and Missouri Deep Loess Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Clayey alluvium  
*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained  
*Slowest permeability:* Very slow (About 0.00 in/hr)  
*Available water capacity:* Moderate (About 6.5 inches)  
*Shrink-swell potential:* Very high (About 17.0 LEP)  
*Flooding hazard:* Rare  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* High  
*Ecological site:* Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 17 inches; silty clay  
 H2—17 to 27 inches; silty clay  
 H3—27 to 60 inches; silty clay

### **MaA—Marshall silty clay loam, 0 to 1 percent slopes**

Map Unit Composition

Marshall: 100 percent

Component Descriptions

Marshall  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Loess hill on upland  
*Parent material:* Fine-silty noncalcareous loess  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.7 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (irrigated):* 1  
*Land capability (nonirrigated):* 1

*Typical Profile:*

H1—0 to 16 inches; silty clay loam  
 H2—16 to 39 inches; silty clay loam  
 H3—39 to 60 inches; silt loam

### **MaB—Marshall silty clay loam, 1 to 3 percent slopes**

Map Unit Composition

Marshall: 100 percent

Component Descriptions

Marshall  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Loess hill on upland  
*Parent material:* Fine-silty noncalcareous loess  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.6 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 14 inches; silty clay loam  
 H2—14 to 38 inches; silty clay loam  
 H3—38 to 60 inches; silt loam

### **MaC—Marshall silty clay loam, 3 to 7 percent slopes**

Map Unit Composition

Marshall: 100 percent

Component Descriptions

Marshall  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Loess hill on upland  
*Parent material:* Fine-silty noncalcareous loess

*Slope:* 3 to 7 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.6 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 12 inches; silty clay loam  
 H2—12 to 37 inches; silty clay loam  
 H3—37 to 60 inches; silt loam

### **MaC2—Marshall silty clay loam, 3 to 7 percent slopes, Eroded**

Map Unit Composition

Marshall: 100 percent

Component Descriptions

Marshall  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Loess hill on upland  
*Parent material:* Fine-silty noncalcareous loess  
*Slope:* 3 to 7 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.5 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 8 inches; silty clay loam  
 H2—8 to 32 inches; silty clay loam

H3—32 to 60 inches; silt loam

### **MaD—Marshall silty clay loam, 7 to 11 percent slopes**

Map Unit Composition

Marshall: 100 percent

Component Descriptions

Marshall  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Loess hill on upland  
*Parent material:* Fine-silty noncalcareous loess  
*Slope:* 7 to 11 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.5 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* High  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 8 inches; silty clay loam  
 H2—8 to 32 inches; silty clay loam  
 H3—32 to 60 inches; silt loam

### **MeD2—Marshall-Ponca silty clay loams, 7 to 11 percent slopes, Eroded**

Map Unit Composition

Marshall: 70 percent  
 Ponca: 30 percent

Component Descriptions

Marshall

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty noncalcareous loess

*Slope:* 7 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 8 inches; silty clay loam

H2—8 to 36 inches; silty clay loam

H3—36 to 60 inches; silt loam

Ponca

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 7 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 5 inches; silty clay loam

H2—5 to 20 inches; silty clay loam

H3—20 to 60 inches; silt loam

## **MfE—Marshall And Ponca Soils, 11 to 17 percent slopes**

Map Unit Composition

Marshall: 50 percent

Ponca: 50 percent

Component Descriptions

Marshall

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty noncalcareous loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 12 inches; silty clay loam

H2—12 to 35 inches; silty clay loam

H3—35 to 60 inches; silt loam

Ponca

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 7 inches; silty clay loam

H2—7 to 22 inches; silty clay loam

H3—22 to 60 inches; silt loam

**MfE2—Marshall And Ponca Soils,  
11 to 17 percent slopes, Eroded**

## Map Unit Composition

Marshall: 50 percent  
Ponca: 50 percent

## Component Descriptions

## Marshall

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty noncalcareous loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 7 inches; silty clay loam

H2—7 to 31 inches; silty clay loam

H3—31 to 60 inches; silt loam

## Ponca

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.3 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 4 inches; silty clay loam

H2—4 to 20 inches; silty clay loam

H3—20 to 60 inches; silt loam

**MoA—Monona silt loam, 0 to 1  
percent slopes**

## Map Unit Composition

Monona: 100 percent

## Component Descriptions

## Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.7 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 1

*Typical Profile:*

H1—0 to 12 inches; silt loam

H2—12 to 35 inches; silt loam

H3—35 to 60 inches; silt loam

**MoB—Monona silt loam, 1 to 3  
percent slopes**

## Map Unit Composition

Monona: 100 percent

## Component Descriptions

## Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 10 inches; silt loam

H2—10 to 33 inches; silt loam

H3—33 to 60 inches; silt loam

## **MoC—Monona silt loam, 3 to 7 percent slopes**

Map Unit Composition

Monona: 100 percent

Component Descriptions

Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 3 to 7 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 10 inches; silt loam

H2—10 to 33 inches; silt loam

H3—33 to 60 inches; silt loam

## **MoC2—Monona silt loam, 3 to 7 percent slopes, Eroded**

Map Unit Composition

Pohocco: 100 percent

Component Descriptions

Pohocco

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 3 to 7 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.7 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 7 inches; silt loam

H2—7 to 30 inches; silty clay loam

H3—30 to 60 inches; silt loam

## **MoD—Monona silt loam, 7 to 11 percent slopes**

Map Unit Composition

Monona: 100 percent

Component Descriptions

Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 7 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 10 inches; silt loam

H2—10 to 33 inches; silt loam

H3—33 to 60 inches; silt loam

## **MoD2—Monona silt loam, 7 to 11 percent slopes, Eroded**

Map Unit Composition

Pohocco: 100 percent

Component Descriptions

Pohocco

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 7 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.7 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 29 inches; silty clay loam

H3—29 to 60 inches; silt loam

## **MoE—Monona silt loam, 11 to 17 percent slopes**

Map Unit Composition

Monona: 100 percent

Component Descriptions

Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 10 inches; silt loam

H2—10 to 33 inches; silt loam

H3—33 to 60 inches; silt loam

## **MsE2—Monona And Ida silt loams, 11 to 17 percent slopes, Eroded**

Map Unit Composition

Pohocco: 50 percent

Ida: 50 percent

Component Descriptions

**Pohocco**

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 11 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 29 inches; silty clay loam

H3—29 to 30 inches; silt loam

**Ida**

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 14 to 17 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland - Veg. Zone 4

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

## **MsF—Monona And Ida silt loams, 17 to 30 percent slopes**

**Map Unit Composition**

Monona: 50 percent

Ida: 50 percent

**Component Descriptions****Monona**

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 17 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 8 inches; silt loam

H2—8 to 31 inches; silt loam

H3—31 to 60 inches; silt loam

**Ida**

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 20 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Limy Upland - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

## **MsF2—Monona And Ida silt loams, 17 to 30 percent slopes, Eroded**



## Map Unit Composition

Monona: 50 percent  
Ida: 50 percent

## Component Descriptions

## Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 17 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 5 inches; silt loam

H2—5 to 27 inches; silt loam

H3—27 to 60 inches; silt loam

## Ida

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 20 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Limy Upland - Veg. Zone 4

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

**MsG—Monona And Ida silt loams,  
30 to 60 percent slopes**

## Map Unit Composition

Monona: 50 percent  
Ida: 50 percent

## Component Descriptions

## Monona

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 30 to 40 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (nonirrigated):* 7e

*Typical Profile:*

H1—0 to 7 inches; silt loam

H2—7 to 30 inches; silt loam

H3—30 to 60 inches; silt loam

## Ida

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess bluff on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 30 to 60 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Thin Loess - Veg. Zone 4

*Land capability (nonirrigated):* 7e

*Typical Profile:*

H1—0 to 5 inches; silt loam  
H2—5 to 60 inches; silt loam

## On—Onawa silty clay

### Map Unit Composition

Onawa: 100 percent

### Component Descriptions

Onawa  
*MLRA*: 107 - Iowa and Missouri Deep Loess Hills  
*Landform*: Flood plain on river valley  
*Parent material*: Calcareous clayey alluvium over loamy alluvium  
*Slope*: 0 to 1 percent  
*Drainage class*: Somewhat poorly drained  
*Slowest permeability*: Very slow (About 0.01 in/hr)  
*Available water capacity*: High (About 9.5 inches)  
*Shrink-swell potential*: Very high (About 12.0 LEP)  
*Flooding hazard*: Occasional  
*Depth to seasonal water saturation*: About 18 to 36 inches  
*Runoff class*: High  
*Ecological site*: Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated)*: 2w  
*Land capability (nonirrigated)*: 2w

### Typical Profile:

H1—0 to 7 inches; silty clay  
H2—7 to 18 inches; silty clay  
H3—18 to 60 inches; very fine sandy loam

### Minor Components

#### Albaton

*Slope*: 0 to 1 percent

## Pa—Percival silty clay

### Map Unit Composition

Percival: 100 percent

### Component Descriptions

Percival

*MLRA*: 107 - Iowa and Missouri Deep Loess Hills

*Landform*: Flood plain on river valley

*Parent material*: Clayey alluvium over sandy alluvium

*Slope*: 0 to 1 percent

*Drainage class*: Somewhat poorly drained

*Slowest permeability*: Very slow (About 0.01 in/hr)

*Available water capacity*: Low (About 4.8 inches)

*Shrink-swell potential*: Very high (About 17.0 LEP)

*Flooding hazard*: Occasional

*Depth to seasonal water saturation*: About 18 to 36 inches

*Runoff class*: High

*Ecological site*: Clayey Overflow - Veg. Zone 4

*Land capability (irrigated)*: 2w

*Land capability (nonirrigated)*: 2w

### Typical Profile:

H1—0 to 7 inches; silty clay  
H2—7 to 16 inches; silty clay  
H3—16 to 60 inches; stratified fine sand to loamy fine sand

## Pb—Pits And Dumps

### Map Unit Composition

Pits: 100 percent

### Component Descriptions

Pits

*MLRA*: 106 - Nebraska and Kansas Loess-Drift Hills

*Slope*: 0 to 30 percent

*Drainage class*: Excessively drained

*Slowest permeability*: Rapid (About 5.95 in/hr)

*Available water capacity*: Low (About 3.5 inches)

*Shrink-swell potential*: Low (About 1.5 LEP)

*Flooding hazard*: None

*Depth to seasonal water saturation*: More than 6 feet

*Runoff class*: Very low

*Land capability (nonirrigated)*: 8s

## Pc—Platte Soils

### Map Unit Composition

Platte: 100 percent

#### Component Descriptions

Platte

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderate (About 0.57 in/hr)

*Available water capacity:* Low (About 3.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated - Veg. Zone 4

*Land capability (irrigated):* 4w

*Land capability (nonirrigated):* 4w

#### Typical Profile:

H1—0 to 9 inches; loam

H2—9 to 12 inches; loam

H3—12 to 60 inches; gravelly coarse sand

#### Minor Components

**Wt At 0-1 Foot**

### **PdD2—Ponca And Ida silt loams, 7 to 11 percent slopes, Eroded**

#### Map Unit Composition

Ponca: 50 percent

Ida: 50 percent

#### Component Descriptions

Ponca

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 7 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.3 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Silty - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

#### Typical Profile:

H1—0 to 4 inches; silt loam

H2—4 to 20 inches; silty clay loam

H3—20 to 60 inches; silt loam

Ida

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty calcareous loess

*Slope:* 9 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland - Veg. Zone 4

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 3e

#### Typical Profile:

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

### **PdE2—Ponca And Ida silt loams, 11 to 17 percent slopes, Eroded**

#### Map Unit Composition

Ponca: 50 percent

Ida: 50 percent

#### Component Descriptions

Ponca

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland

*Parent material:* Fine-silty loess

*Slope:* 11 to 17 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.3 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* High  
*Ecological site:* Silty - Veg. Zone 4  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 4 inches; silt loam  
 H2—4 to 20 inches; silty clay loam  
 H3—20 to 60 inches; silt loam

**Ida**

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills

*Landform:* Loess hill on upland  
*Parent material:* Fine-silty calcareous loess  
*Slope:* 14 to 17 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.4 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Limy Upland - Veg. Zone 4  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 5 inches; silt loam  
 H2—5 to 60 inches; silt loam

## **Ra—Riverwash**

**Map Unit Composition**

Riverwash: 100 percent

**Component Descriptions**

Riverwash  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills

*Landform:* Flood plain on river valley  
*Parent material:* Stratified sandy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Very low (About 2.6 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Frequent  
*Depth to seasonal water saturation:* About 0 to 0 inches  
*Runoff class:* Negligible  
*Land capability (nonirrigated):* 8w

*Typical Profile:*

H1—0 to 6 inches; sand  
 H2—6 to 60 inches; stratified sand to very gravelly coarse sand

## **Rk—Rock Land**

**Map Unit Composition**

Rock outcrop: 100 percent

**Component Descriptions**

Rock outcrop  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Bluff on upland  
*Parent material:* Residuum weathered from limestone and sandstone  
*Slope:* 9 to 60 percent  
*Depth to restrictive feature:* 0 inches to bedrock (paralithic)  
*Drainage class:* Somewhat excessively drained  
*Available water capacity:* Very low (About 0.0 inches)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very high  
*Ecological site:* No Site - Veg. Zone 4  
*Land capability (nonirrigated):* 8s

## **Rn—Rough Broken Land, Loess**

**Map Unit Composition**

Ida Variant: 100 percent

**Component Descriptions**

Ida Variant

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills*Landform:* Gully on loess hill on upland*Parent material:* Fine-silty loess*Slope:* 30 to 60 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 11.3 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* High*Ecological site:* Thin Loess - Veg. Zone 4*Land capability (nonirrigated):* 7e**Typical Profile:**

H1—0 to 6 inches; silt loam

H2—6 to 60 inches; silt loam

**Sd—sandy Alluvial Land****Map Unit Composition**

Inavale: 100 percent

**Component Descriptions**

Inavale

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills*Landform:* Flood plain on river valley*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.7 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Frequent*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 4*Land capability (nonirrigated):* 6w**Typical Profile:**

H1—0 to 6 inches; sand

H2—6 to 60 inches; fine sand

**Minor Components****Wt At 0-1 Foot****Sp—Sarpy fine sand****Map Unit Composition**

Sarpy: 100 percent

**Component Descriptions**

Sarpy

*MLRA:* 107 - Iowa and Missouri Deep Loess Hills*Landform:* Flood plain on river valley*Parent material:* Calcareous sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.2 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 4*Land capability (irrigated):* 4s*Land capability (nonirrigated):* 6s**Typical Profile:**

H1—0 to 6 inches; fine sand

H2—6 to 60 inches; fine sand

**Ss—silty Alluvial Land****Map Unit Composition**

Nodaway Variant: 100 percent

**Component Descriptions**

Nodaway Variant

*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills*Landform:* Flood plain on river valley*Parent material:* Silty alluvium*Slope:* 0 to 2 percent*Drainage class:* Moderately well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 11.4 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Frequent  
*Depth to seasonal water saturation:* About 36 to 72 inches  
*Runoff class:* Low  
*Ecological site:* Silty Overflow - Veg. Zone 4  
*Land capability (nonirrigated):* 6w

*Typical Profile:*  
 H1—0 to 7 inches; silt loam  
 H2—7 to 80 inches; silt loam

**Minor Components**  
**Ponded Soils**

**StE2—Steinauer clay loam, 11 to 30 percent slopes, Eroded**

Map Unit Composition

Steinauer: 100 percent

Component Descriptions  
 Steinauer  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Hill on upland  
*Parent material:* Calcareous loamy till  
*Slope:* 11 to 30 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 10.8 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very high  
*Ecological site:* Limey Upland - Veg. Zone 4  
*Land capability (nonirrigated):* 6e

*Typical Profile:*  
 H1—0 to 6 inches; clay loam  
 H2—6 to 18 inches; clay loam  
 H3—18 to 60 inches; clay loam

**Wb—Wabash silt loam**

Map Unit Composition

Wabash: 100 percent

Component Descriptions  
 Wabash  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Clayey alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Very slow (About 0.00 in/hr)  
*Available water capacity:* Moderate (About 7.9 inches)  
*Shrink-swell potential:* Very high (About 17.0 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* High  
*Ecological site:* Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*  
 H1—0 to 15 inches; silt loam  
 H2—15 to 60 inches; silty clay

**Wc—Wabash silty clay**

Map Unit Composition

Wabash: 100 percent

Component Descriptions  
 Wabash  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Clayey alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Very slow (About 0.00 in/hr)  
*Available water capacity:* Moderate (About 6.1 inches)  
*Shrink-swell potential:* Very high (About 17.0 LEP)

*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* High  
*Ecological site:* Clayey Overflow - Veg. Zone 4  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*  
 H1—0 to 7 inches; silty clay  
 H2—7 to 60 inches; silty clay

## **Wm—Wann fine sandy loam**

Map Unit Composition

Wann: 100 percent

Component Descriptions

Wann  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Calcareous loamy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Moderate (About 9.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 18 to 42 inches  
*Runoff class:* Negligible  
*Ecological site:* Subirrigated - Veg. Zone 4  
*Land capability (irrigated):* 2w  
*Land capability (nonirrigated):* 2w

*Typical Profile:*  
 H1—0 to 23 inches; fine sandy loam  
 H2—23 to 50 inches; fine sandy loam  
 H3—50 to 60 inches; stratified sandy clay loam to fine sand

**Minor Components**  
**Ponded Soils**

## **Wt—Wet Alluvial Land**

Map Unit Composition

Barney: 100 percent

Component Descriptions

Barney  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Low (About 4.2 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Frequent  
*Depth to seasonal water saturation:* About 0 to 12 inches  
*Runoff class:* Negligible  
*Ecological site:* Wet Land - Veg. Zone 4  
*Land capability (nonirrigated):* 5w  
  
*Typical Profile:*  
 H1—0 to 7 inches; loam  
 H2—7 to 15 inches; stratified loam to sand  
 H3—15 to 60 inches; gravelly coarse sand

## **zwa—Water < 40 Acres**

Map Unit Composition

Water: 100 percent

Component Descriptions

Water  
*MLRA:* 106 - Nebraska and Kansas Loess-Drift Hills  
*Depth to seasonal water saturation:* More than 6 feet

## **zwb—Water > 40 Acres**

Map Unit Composition

Water: 100 percent

## Component Descriptions

Water

*MLRA*: 106 - Nebraska and Kansas Loess-Drift  
Hills*Depth to seasonal water saturation*: More than 6  
feet



Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive land-forming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes. In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

(Class 1) soils have slight limitations that restrict their use.

(Class 2) soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

(Class 3) soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

(Class 4) soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

(Class 5) soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 6) soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 7) soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

(Class 8) soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, e, w, s, or c, to the class numeral, for example, 2e. The letter e shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; w shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); s shows that the soil is limited mainly because it is shallow, droughty, or stony; and c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by w, s, or c because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

The capability classification of map units in this survey area is given in the section "Detailed Soil Map Units" and in the Land Capability and Component Yields table.

#### Crop Yield Estimates

The average yields per acre that can be expected of the principal crops under a high level of management are shown in "Land Capability and Component Yields" table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, animal waste manure, and green manure crops; and harvesting that ensures the smallest possible loss.

For yields of irrigated crops, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in this table, are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service (NRCS) or the Cooperative Extension Service (CES) can provide information about the management and productivity of the soils for those crops.

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 2 of 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Soybeans		Winter wheat	
	N	I	N	I	N	I	N	I	N	I
			Bu		Bu		Bu		Bu	
Ab: ALBATON-----	3w	---	96.00	---	75.00	---	32.00	---	35.00	---
Ac: ALBATON-----	3w	3w	90.00	---	70.00	---	30.00	---	30.00	---
Af: ALDA-----	3w	3w	72.00	---	65.00	---	30.00	---	30.00	---
Ag: ALDA-----	3w	3w	96.00	---	85.00	---	35.00	---	30.00	---
Ca: CARR-----	2w	2w	108.00	---	70.00	---	30.00	---	30.00	---
Cc: CASS-----	2e	2e	102.00	---	65.00	---	30.00	---	30.00	---
Cd: CASS-----	2e	2e	120.00	---	95.00	---	35.00	---	35.00	---
Ce: CASS-----	1	1	126.00	---	95.00	---	35.00	---	35.00	---
Cg: COLO-----	2w	2w	120.00	---	105.00	---	30.00	---	35.00	---
Ck: COLO-----	2w	2w	144.00	---	110.00	---	30.00	---	35.00	---
KENNEBEC-----	2w	---	144.00	---	110.00	---	30.00	---	35.00	---
Cm: URBAN LAND-----	---	---	---	---	---	---	---	---	---	---
DcE: DICKINSON-----	6e	---	84.00	---	---	---	20.00	---	20.00	---
Ed: EUDORA-----	1	1	144.00	---	120.00	---	45.00	---	40.00	---
Ga: GIBBON-----	3w	3w	96.00	---	75.00	---	25.00	---	30.00	---
Gb: GIBBON-----	2w	2w	150.00	---	115.00	---	45.00	---	40.00	---
Gc: GIBBON-----	2w	2w	144.00	---	110.00	---	35.00	---	35.00	---
Gs: GIBBON-----	2w	2w	48.00	---	---	---	---	---	25.00	---
SALTINE-----	6s	---	48.00	---	---	---	---	---	25.00	---
Gu: IDA VARIANT-----	7e	---	---	---	25.00	---	25.00	---	---	---
Ha: HAYNIE-----	2w	2w	120.00	---	100.00	---	40.00	---	40.00	---
IdD2: IDA-----	4e	---	60.00	---	50.00	---	---	---	30.00	---
IdE: IDA-----	6e	---	---	---	---	---	---	---	---	---
IdE2: IDA-----	6e	---	---	---	20.00	---	20.00	---	---	---
Im: INAVALE-----	4e	3e	54.00	---	45.00	---	---	---	20.00	---
In: INAVALE-----	6e	4e	34.00	---	---	---	---	---	---	---
JuB: JUDSON-----	2e	3e	126.00	---	95.00	---	42.00	---	40.00	---

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 3 of 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Soybeans		Winter wheat	
	N	I	N	I	N	I	N	I	N	I
			Bu		Bu		Bu		Bu	
Ke: KENNEBEC-----	2w	2w	132.00	---	105.00	---	45.00	---	35.00	---
Le: LOCKTON-----	3w	3w	132.00	---	110.00	---	40.00	---	35.00	---
Ls: LUTON-----	3w	3w	102.00	---	90.00	---	35.00	---	35.00	---
Lt: LUTON-----	3w	3w	96.00	---	80.00	---	32.00	---	32.00	---
Lu: LUTON-----	3w	3w	90.00	---	75.00	---	30.00	---	30.00	---
MaA: MARSHALL-----	1	1	114.00	---	100.00	---	40.00	---	40.00	---
MaB: MARSHALL-----	2e	2e	114.00	---	95.00	---	40.00	---	40.00	---
MaC: MARSHALL-----	2e	3e	108.00	---	90.00	---	35.00	---	38.00	---
MaC2: MARSHALL-----	3e	2e	102.00	---	85.00	---	30.00	---	35.00	---
MaD: MARSHALL-----	3e	4e	96.00	---	80.00	---	28.00	---	35.00	---
MeD2: MARSHALL-----	3e	4e	90.00	---	75.00	---	26.00	---	32.00	---
PONCA-----	3e	4e	90.00	---	75.00	---	25.00	---	32.00	---
MfE: MARSHALL-----	4e	---	84.00	---	70.00	---	24.00	---	30.00	---
PONCA-----	4e	---	84.00	---	70.00	---	---	---	30.00	---
MfE2: MARSHALL-----	4e	---	78.00	---	65.00	---	---	---	28.00	---
PONCA-----	4e	---	78.00	---	65.00	---	---	---	28.00	---
MoA: MONONA-----	1	1	114.00	---	100.00	---	40.00	---	40.00	---
MoB: MONONA-----	2e	2e	114.00	---	95.00	---	40.00	---	40.00	---
MoC: MONONA-----	2e	3e	108.00	---	90.00	---	35.00	---	38.00	---
MoC2: POHOCCO-----	2e	3e	102.00	---	85.00	---	30.00	---	35.00	---
MoD: MONONA-----	3e	4e	96.00	---	85.00	---	28.00	---	30.00	---
MoD2: POHOCCO-----	3e	4e	90.00	---	80.00	---	26.00	---	28.00	---
MoE: MONONA-----	4e	---	84.00	---	75.00	---	24.00	---	25.00	---
MsE2: IDA-----	4e	---	78.00	---	75.00	---	---	---	20.00	---
POHOCCO-----	4e	---	78.00	---	75.00	---	23.00	---	20.00	---
MsF: IDA-----	6e	---	66.00	---	---	---	22.00	---	24.00	---
MONONA-----	6e	---	66.00	---	---	---	22.00	---	24.00	---
MsF2: IDA-----	6e	---	---	---	---	---	---	---	---	---
MONONA-----	6e	---	---	---	---	---	---	---	---	---

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 4 of 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Corn		Grain sorghum		Soybeans		Winter wheat	
	N	I	N	I	N	I	N	I	N	I
			Bu		Bu		Bu		Bu	
MsG:										
IDA-----	7e	---	54.00	---	---	---	20.00	---	22.00	---
MONONA-----	7e	---	54.00	---	---	---	20.00	---	22.00	---
On:										
ONAWA-----	2w	2w	108.00	---	85.00	---	35.00	---	40.00	---
Pa:										
PERCIVAL-----	2w	2w	90.00	---	75.00	---	30.00	---	30.00	---
Pb:										
PITS-----	8s	---	---	---	---	---	---	---	---	---
Pc:										
PLATTE-----	4w	4w	36.00	---	30.00	---	---	---	15.00	---
PdD2:										
IDA-----	3e	4e	78.00	---	80.00	---	---	---	28.00	---
PONCA-----	3e	4e	78.00	---	80.00	---	---	---	28.00	---
PdE2:										
IDA-----	4e	---	72.00	---	65.00	---	---	---	20.00	---
PONCA-----	4e	---	72.00	---	65.00	---	---	---	20.00	---
Ra:										
RIVERWASH-----	8w	---	---	---	---	---	---	---	---	---
Rk:										
ROCK OUTCROP----	8s	---	---	---	---	---	---	---	---	---
Rn:										
IDA VARIANT-----	7e	---	54.00	---	---	---	22.00	---	22.00	---
Sd:										
INAVALE-----	6w	---	---	---	18.00	---	28.00	---	---	---
Sp:										
SARPY-----	6s	4s	---	---	---	---	---	---	---	---
Ss:										
NODAWAY VARIANT--	6w	---	---	---	---	---	---	---	---	---
StE2:										
STEINAUER-----	6e	---	54.00	---	50.00	---	18.00	---	24.00	---
Wb:										
WABASH-----	3w	3w	96.00	---	85.00	---	35.00	---	32.00	---
Wc:										
WABASH-----	3w	3w	90.00	---	60.00	---	30.00	---	30.00	---
Wm:										
WANN-----	2w	2w	84.00	---	70.00	---	35.00	---	30.00	---
Wt:										
BARNEY-----	5w	---	---	---	---	---	---	---	---	---
zwa:										
WATER-----	---	---	---	---	---	---	---	---	---	---
zwb:										
WATER-----	---	---	---	---	---	---	---	---	---	---



Farmland Classification  
Douglas and Sarpy Counties, Nebraska : Out-of-date

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
Ca	Carr fine sandy loam	All areas are prime farmland
Cc	Cass fine sandy loam	All areas are prime farmland
Cd	Cass fine sandy loam, loamy substratum	All areas are prime farmland
Ce	Cass very fine sandy loam	All areas are prime farmland
Ed	Eudora silt loam	All areas are prime farmland
Ha	Haynie silt loam	All areas are prime farmland
JuB	Judson silt loam, 3 to 7 percent slopes	All areas are prime farmland
Ke	Kennebec silt loam, occasionally flooded	All areas are prime farmland
MaA	Marshall silty clay loam, 0 to 1 percent slopes	All areas are prime farmland
MaB	Marshall silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
MaC	Marshall silty clay loam, 3 to 7 percent slopes	All areas are prime farmland
MaC2	Marshall silty clay loam, 3 to 7 percent slopes, eroded	All areas are prime farmland
MoA	Monona silt loam, 0 to 1 percent slopes	All areas are prime farmland
MoB	Monona silt loam, 1 to 3 percent slopes	All areas are prime farmland
MoC	Monona silt loam, 3 to 7 percent slopes	All areas are prime farmland
MoC2	Monona silt loam, 3 to 7 percent slopes, eroded	All areas are prime farmland
Af	Alda fine sandy loam	Prime farmland if drained
Ag	Alda very fine sandy loam	Prime farmland if drained
Cg	Colo silty clay loam	Prime farmland if drained
Ck	Colo and kennebec soils	Prime farmland if drained
Ga	Gibbon loamy sand, overwash	Prime farmland if drained
Gb	Gibbon silt loam	Prime farmland if drained
Gc	Gibbon silty clay loam	Prime farmland if drained
Le	Lex soils, noncalcareous variant	Prime farmland if drained
On	Onawa silty clay	Prime farmland if drained
Pa	Percival silty clay	Prime farmland if drained
Wm	Wann fine sandy loam	Prime farmland if drained



The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
Ab	Albaton Silt Loam, Overwash-----	35
Ac	Albaton Silty Clay-----	23
Af	Alda Fine Sandy Loam-----	26
Ag	Alda Very Fine Sandy Loam-----	27
Ca	Carr Fine Sandy Loam-----	51
Cc	Cass Fine Sandy Loam-----	54
Cd	Cass Fine Sandy Loam, Loamy Substratum-----	69
Ce	Cass Very Fine Sandy Loam-----	55
Cg	Colo Silty Clay Loam-----	59
Ck	Colo And Kennebec Soils-----	66
Cm	Cut And Fill Land-----	15
DcE	Dickinson Soils, 11 To 17 Percent Slopes-----	44
Ed	Eudora Silt Loam-----	70
Ga	Gibbon Loamy Sand, Overwash-----	51
Gb	Gibbon Silt Loam-----	54
Gc	Gibbon Silty Clay Loam-----	54
Gs	Gibbon-Slickspots Complex-----	43
Gu	Gullied Land-----	3
Ha	Haynie Silt Loam-----	48
IdD2	Ida Silt Loam, 7 To 17 Percent Slopes, Eroded-----	40
IdE	Ida Silt Loam, 17 To 30 Percent Slopes-----	18
IdE2	Ida Silt Loam, 17 To 30 Percent Slopes, Eroded-----	18
Im	Inavale Loamy Fine Sand-----	28
In	Inavale Loamy Fine Sand, Hummocky-----	25
JuB	Judson Silt Loam, 3 To 7 Percent Slopes-----	73
Ke	Kennebec Silt Loam, Occasionally Flooded-----	73
Le	Lex Soils, Noncalcareous Variant-----	51
Ls	Luton Silt Loam, Overwash-----	43
Lt	Luton Silty Clay Loam-----	41
Lu	Luton Silty Clay-----	35
MaA	Marshall Silty Clay Loam, 0 To 1 Percent Slopes-----	79
MaB	Marshall Silty Clay Loam, 1 To 3 Percent Slopes-----	77
MaC	Marshall Silty Clay Loam, 3 To 7 Percent Slopes-----	74
MaC2	Marshall Silty Clay Loam, 3 To 7 Percent Slopes, Eroded-----	74
MaD	Marshall Silty Clay Loam, 7 To 11 Percent Slopes-----	69
MeD2	Marshall-Ponca Silty Clay Loams, 7 To 11 Percent Slopes, Eroded-----	67
MfE	Marshall And Ponca Soils, 11 To 17 Percent Slopes-----	60
MfE2	Marshall And Ponca Soils, 11 To 17 Percent Slopes, Eroded-----	59
MoA	Monona Silt Loam, 0 To 1 Percent Slopes-----	74
MoB	Monona Silt Loam, 1 To 3 Percent Slopes-----	73
MoC	Monona Silt Loam, 3 To 7 Percent Slopes-----	70
MoC2	Monona Silt Loam, 3 To 7 Percent Slopes, Eroded-----	70
MoD	Monona Silt Loam, 7 To 11 Percent Slopes-----	65
MoD2	Monona Silt Loam, 7 To 11 Percent Slopes, Eroded-----	65
MoE	Monona Silt Loam, 11 To 17 Percent Slopes-----	58
MsE2	Monona And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded-----	40
MsF	Monona And Ida Silt Loams, 17 To 30 Percent Slopes-----	22
MsF2	Monona And Ida Silt Loams, 17 To 30 Percent Slopes, Eroded-----	22
MsG	Monona And Ida Silt Loams, 30 To 60 Percent Slopes-----	5
On	Onawa Silty Clay-----	36
Pa	Percival Silty Clay-----	25
Pb	Pits And Dumps-----	18
Pc	Platte Soils-----	33
PdD2	Ponca And Ida Silt Loams, 7 To 11 Percent Slopes, Eroded-----	54
PdE2	Ponca And Ida Silt Loams, 11 To 17 Percent Slopes, Eroded-----	47
Ra	Riverwash-----	0
Rk	Rock Land-----	0
Rn	Rough Broken Land, Loess-----	3
Sd	Sandy Alluvial Land-----	22
Sp	Sarpy Fine Sand-----	22
Ss	Silty Alluvial Land-----	56
StE2	Steinauer Clay Loam, 11 To 30 Percent Slopes, Eroded-----	27
Wb	Wabash Silt Loam-----	50
Wc	Wabash Silty Clay-----	41
Wm	Wann Fine Sandy Loam-----	47
Wt	Wet Alluvial Land-----	24
zwa	Water < 40 Acres-----	0
zwb	Water > 40 Acres-----	0





Douglas and Sarpy Counties, Nebraska: Out-of-date  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Ab:ALBATON-----	100	N/A	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.37	.37	5	4L	86
Ac:ALBATON-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.28	.28	5	4	86
Af:ALDA-----	100	3w-	3w	Prime farmland if drained	C	Subirrigated - Veg. Zone 4		.20	.20	4	3	86
Ag:ALDA-----	100	3w-	3w	Prime farmland if drained	C	Subirrigated - Veg. Zone 4		.28	.28	4	3	86
Ca:CARR-----	100	2w-	2w	All areas are prime farmland	B	Sandy Lowland - Veg. Zone 4		.24	.24	4	3	86
Cc:CASS-----	100	2e-	2e	All areas are prime farmland	B	Sandy Lowland - Veg. Zone 4		.20	.20	5	3	86
Cd:CASS-----	100	2e-	2e	All areas are prime farmland	B	Sandy Lowland - Veg. Zone 4		.20	.20	5	3	86
Ce:CASS-----	100	1-	1	All areas are prime farmland	B	Sandy Lowland - Veg. Zone 4		.32	.32	5	3	86
Cg:COLO-----	100	2w-	2w	Prime farmland if drained	B/D	Subirrigated - Veg. Zone 4		.28	.28	5	7	38
Ck:COLO-----	50	2w-	2w	Prime farmland if drained	B/D	Subirrigated - Veg. Zone 4		.28	.28	5	7	38
Ck:KENNEBEC-----	50	N/A	2w	Prime farmland if drained	B	Silty Lowland - Veg. Zone 4		.28	.28	5	6	48
Cm:URBAN LAND---	100	N/A	N/A	Not prime farmland	D	Unspecified		---	---	-	8	0
DcE:DICKINSON---	100	N/A	6e	Not prime farmland	B	Sandy - Veg. Zone 4		.20	.20	4	3	86
Ed:EUDORA-----	100	1-	1	All areas are prime farmland	B	Silty Overflow - Veg. Zone 4		.32	.32	5	5	56
Ga:GIBBON-----	100	3w-	3w	Prime farmland if drained	B	Subirrigated - Veg. Zone 4		.17	.17	4	2	134
Gb:GIBBON-----	100	2w-	2w	Prime farmland if drained	B	Subirrigated - Veg. Zone 4		.32	.32	5	4L	86
Gc:GIBBON-----	100	2w-	2w	Prime farmland if drained	B	Subirrigated - Veg. Zone 4		.32	.32	5	4L	86

Douglas and Sarpy Counties, Nebraska: Out-of-date  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Gs:GIBBON-----	65	2w-	2w	Not prime farmland	B	Subirrigated - Veg. Zone 4		.32	.32	5	4L	86
Gs:SALTINE-----	35	N/A	6s	Not prime farmland	C	Saline Subirrigated - Veg. Zone 4		.37	.37	5	4L	86
Gu:IDA VARIANT--	100	N/A	7e	Not prime farmland	B	Thin Loess - Veg. Zone 4		.43	.43	4	4L	86
Ha:HAYNIE-----	100	2w-	2w	All areas are prime farmland	B	Silty Lowland - Veg. Zone 4		.32	.32	5	4L	86
IdD2:IDA-----	100	N/A	4e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
IdE:IDA-----	100	N/A	6e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.32	.32	5	4L	86
IdE2:IDA-----	100	N/A	6e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
Im:INAVALE-----	100	3e-	4e	Not prime farmland	A	Sandy Lowland - Veg. Zone 4		.17	.17	5	2	134
In:INAVALE-----	100	4e-	6e	Not prime farmland	A	Sandy Lowland - Veg. Zone 4		.17	.17	5	2	134
JuB:JUDSON-----	100	3e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
Ke:KENNEBEC-----	100	2w-	2w	All areas are prime farmland	B	Silty Overflow - Veg. Zone 4		.28	.28	5	6	48
Le:LOCKTON-----	100	3w-	3w	Prime farmland if drained	B	Subirrigated - Veg. Zone 4		.28	.28	4	5	56
Ls:LUTON-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.43	.43	5	6	48
Lt:LUTON-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.32	.32	5	4	86
Lu:LUTON-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.28	.28	5	4	86
MaA:MARSHALL----	100	1-	1	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	7	38
MaB:MARSHALL----	100	2e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	7	38
MaC:MARSHALL----	100	3e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	7	38

Douglas and Sarpy Counties, Nebraska: Out-of-date  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
MaC2:MARSHALL---	100	2e-	3e	All areas are prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MaD:MARSHALL----	100	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	7	38
MeD2:MARSHALL---	70	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MeD2:PONCA-----	30	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MfE:MARSHALL----	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	7	38
MfE:PONCA-----	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MfE2:MARSHALL---	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MfE2:PONCA-----	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	7	38
MoA:MONONA-----	100	1-	1	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MoB:MONONA-----	100	2e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MoC:MONONA-----	100	3e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MoC2:POHOCCO----	100	3e-	2e	All areas are prime farmland	B	Silty - Veg. Zone 4		.32	.37	5	6	48
MoD:MONONA-----	100	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MoD2:POHOCCO----	100	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.37	5	6	48
MoE:MONONA-----	100	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MsE2:POHOCCO----	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.37	5	6	48
MsE2:IDA-----	50	N/A	4e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
MsF:MONONA-----	50	N/A	6e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MsF:IDA-----	50	N/A	6e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.32	.32	5	4L	86

Douglas and Sarpy Counties, Nebraska: Out-of-date  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
MsF2:MONONA-----	50	N/A	6e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MsF2:IDA-----	50	N/A	6e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
MsG:MONONA-----	50	N/A	7e	Not prime farmland	B	Silty - Veg. Zone 4		.28	.28	5	6	48
MsG:IDA-----	50	N/A	7e	Not prime farmland	B	Thin Loess - Veg. Zone 4		.32	.32	5	4L	86
On:ONAWA-----	100	2w-	2w	Prime farmland if drained	D	Clayey Overflow - Veg. Zone 4		.32	.32	5	4	86
Pa:PERCIVAL-----	100	2w-	2w	Prime farmland if drained	C	Clayey Overflow - Veg. Zone 4		.28	.28	4	4	86
Pb:PITS-----	100	N/A	8s	Not prime farmland	A	Unspecified		.10	.17	2	8	0
Pc:PLATTE-----	100	4w-	4w	Not prime farmland	B	Subirrigated - Veg. Zone 4		.28	.28	5	4L	86
PdD2:PONCA-----	50	4e-	3e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	6	48
PdD2:IDA-----	50	4e-	3e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
PdE2:PONCA-----	50	N/A	4e	Not prime farmland	B	Silty - Veg. Zone 4		.32	.32	5	6	48
PdE2:IDA-----	50	N/A	4e	Not prime farmland	B	Limy Upland - Veg. Zone 4		.43	.43	4	4L	86
Ra:RIVERWASH----	100	N/A	8w	Not prime farmland	A	Unspecified		.15	.15	2	1	220
Rk:ROCK OUTCROP-	100	N/A	8s	Not prime farmland	D	No Site - Veg. Zone 4		---	---	1	8	0
Rn:IDA VARIANT--	100	N/A	7e	Not prime farmland	B	Thin Loess - Veg. Zone 4		.43	.43	5	4L	86
Sd:INVALE-----	100	N/A	6w	Not prime farmland	A	Sandy Lowland - Veg. Zone 4		.15	.15	5	1	220
Sp:SARPY-----	100	4s-	6s	Not prime farmland	A	Sandy Lowland - Veg. Zone 4		.15	.15	5	1	220
Ss:NODAWAY VARIANT-----	100	N/A	6w	Not prime farmland	B	Silty Overflow - Veg. Zone 4		.32	.32	5	6	48
StE2:STEINAUER--	100	N/A	6e	Not prime farmland	B	Limey Upland - Veg. Zone 4		.32	.32	5	4L	86
Wb:WABASH-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.28	.28	5	6	48

Douglas and Sarpy Counties, Nebraska: Out-of-date  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Wc:WABASH-----	100	3w-	3w	Not prime farmland	D	Clayey Overflow - Veg. Zone 4		.28	.28	5	4	86
Wm:WANN-----	100	2w-	2w	Prime farmland if drained	B	Subirrigated - Veg. Zone 4		.20	.20	5	3	86
Wt:BARNEY-----	100	N/A	5w	Not prime farmland	D	Wet Land - Veg. Zone 4		.28	.28	5	8	0
zwa:WATER-----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---
zwb:WATER-----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---



RANGELAND PRODUCTIVITY  
Douglas and Sarpy Counties,  
Nebraska

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

**Rangeland.** Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

**Grazed Forest Land.** Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

**Native Pasture.** Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

**Rangeland**

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.



RANGELAND PRODUCTIVITY--Continued  
Douglas and Sarpy Counties,  
Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Ab:				
Albaton-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
Ac:				
Albaton-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
Af:				
Alda-----	Subirrigated - Veg. Zone 4	5,900	5,500	5,100
Ag:				
Alda-----	Subirrigated - Veg. Zone 4	5,900	5,500	5,100
Ca:				
Carr-----	Sandy Lowland - Veg. Zone 4	6,500	5,500	4,500
Cc:				
Cass-----	Sandy Lowland - Veg. Zone 4	4,300	4,000	3,700
Cd:				
Cass-----	Sandy Lowland - Veg. Zone 4	5,000	4,000	2,500
Ce:				
Cass-----	Sandy Lowland - Veg. Zone 4	5,300	4,900	4,500
Cg:				
Colo-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Ck:				
Colo-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Kennebec-----	Silty Lowland - Veg. Zone 4	5,300	4,900	4,500
Cm:				
Urban Land-----	---	---	---	---
DcE:				
Dickinson-----	Sandy - Veg. Zone 4	---	---	---
Ed:				
Eudora-----	Silty Overflow - Veg. Zone 4	10,000	8,000	6,000
Ga:				
Gibbon-----	Subirrigated - Veg. Zone 4	5,000	4,500	3,500
Gb:				
Gibbon-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Gc:				
Gibbon-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Gs:				
Gibbon-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Saltine-----	Saline Subirrigated - Veg. Zone 4	4,300	3,900	3,500
Gu:				
Ida Variant-----	Thin Loess - Veg. Zone 4	2,500	2,300	2,000
Ha:				
Haynie-----	Silty Lowland - Veg. Zone 4	5,300	4,900	4,500
IdD2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
IdE:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
IdE2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Im:				
Inavale-----	Sandy Lowland - Veg. Zone 4	4,300	3,500	2,700
In:				
Inavale-----	Sandy Lowland - Veg. Zone 4	4,000	3,300	2,500
JuB:				
Judson-----	Silty - Veg. Zone 4	4,800	3,900	3,000
Ke:				
Kennebec-----	Silty Overflow - Veg. Zone 4	5,300	4,900	4,500
Le:				
Lockton-----	Subirrigated - Veg. Zone 4	5,200	4,900	4,600
Ls:				
Luton-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
Lt:				
Luton-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
Lu:				
Luton-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
MaA:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MaB:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MaC:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MaC2:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MaD:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MeD2:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
Ponca-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MfE:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
Ponca-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MfE2:				
Marshall-----	Silty - Veg. Zone 4	4,800	4,400	4,000
Ponca-----	Silty - Veg. Zone 4	4,800	4,400	4,000
MoA:				
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200

RANGELAND PRODUCTIVITY--Continued  
Douglas and Sarpy Counties,  
Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
MoB:				
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MoC:				
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MoC2:				
Pohocco-----	Silty - Veg. Zone 4	4,200	3,600	3,000
MoD:				
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MoD2:				
Pohocco-----	Silty - Veg. Zone 4	4,200	3,600	3,000
MoE:				
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MsE2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Pohocco-----	Silty - Veg. Zone 4	4,200	3,600	3,000
MsF:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MsF2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
MsG:				
Ida-----	Thin Loess - Veg. Zone 4	3,500	3,300	3,000
Monona-----	Silty - Veg. Zone 4	4,000	3,600	3,200
On:				
Onawa-----	Clayey Overflow - Veg. Zone 4	4,000	3,600	3,200
Pa:				
Percival-----	Clayey Overflow - Veg. Zone 4	3,800	3,300	2,800
Pb:				
Pits-----	---	---	---	---
Pc:				
Platte-----	Subirrigated - Veg. Zone 4	5,500	5,100	4,700
PdD2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Ponca-----	Silty - Veg. Zone 4	4,800	4,400	4,000
PdE2:				
Ida-----	Limy Upland - Veg. Zone 4	4,000	3,600	3,200
Ponca-----	Silty - Veg. Zone 4	4,800	4,400	4,000
Ra:				
Riverwash-----	---	---	---	---
Rk:				
Rock Outcrop-----	No Site - Veg. Zone 4	0	0	0
Rn:				
Ida Variant-----	Thin Loess - Veg. Zone 4	2,500	2,300	2,000
Sd:				
Inavale-----	Sandy Lowland - Veg. Zone 4	4,300	3,500	2,700
Sp:				
Sarpy-----	Sandy Lowland - Veg. Zone 4	3,800	3,500	3,000
Ss:				
Nodaway Variant-----	Silty Overflow - Veg. Zone 4	3,000	2,700	2,500
StE2:				
Steinauer-----	Limey Upland - Veg. Zone 4	4,000	3,100	2,500
Wb:				
Wabash-----	Clayey Overflow - Veg. Zone 4	10,000	9,000	7,000
Wc:				
Wabash-----	Clayey Overflow - Veg. Zone 4	10,000	9,000	7,000
Wm:				
Wann-----	Subirrigated - Veg. Zone 4	6,300	5,900	5,500
Wt:				
Barney-----	Wet Land - Veg. Zone 4	6,500	6,000	5,500
zwa:				
Water-----	---	---	---	---
zwb:				
Water-----	---	---	---	---



BUILDING SITE DEVELOPMENT  
Douglas and Sarpy Counties, Nebraska

### Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. These tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00  1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00  1.00
Ac: Albaton-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00  1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00  1.00
Af: Alda-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Ag: Alda-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Ca: Carr-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Cc: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Cd: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.95 0.50	Very limited Flooding	1.00
Ce: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Cg: Colo-----	100	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
Ck: Colo-----	50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
Kennebec-----	50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 0.50 0.35	Very limited Flooding Shrink-swell	1.00 0.50
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
Ed: Eudora-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ga: Gibbon-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Gb: Gibbon-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Gc: Gibbon-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50  0.07	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50  0.07
Gs: Gibbon-----	65	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50 0.07	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50 0.07
Saltine-----	35	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50 0.07	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 0.50 0.07
Gu: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ha: Haynie-----	100	Not limited		Not limited		Not limited	
IdD2: Ida-----	100	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
IdE: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
IdE2: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Im: Inavale-----	100	Not limited		Not limited		Not limited	
In: Inavale-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
JuB: Judson-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
Ke: Kennebec-----	100	Very limited Flooding	1.00	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 0.50 0.35	Very limited Flooding	1.00
Le: Lockton-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Ls: Luton-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00
Lt: Luton-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00
Lu: Luton-----	100	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell  Depth to saturated zone	1.00 1.00 1.00
MaA: Marshall-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MaB: Marshall-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
MaC: Marshall-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
MaC2: Marshall-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
MaD: Marshall-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Shrink-swell Slope	0.50 0.04	Very limited Slope Shrink-swell	1.00 0.50
MeD2: Marshall-----	70	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Shrink-swell Slope	0.50 0.04	Very limited Slope Shrink-swell	1.00 0.50
Ponca-----	30	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
MfE: Marshall-----	50	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Slope Shrink-swell	1.00 0.50
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
MfE2: Marshall-----	50	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Slope Shrink-swell	1.00 0.50
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
MoA: Monona-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
MoB: Monona-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
MoC: Monona-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell Slope	0.50 0.12
MoC2: Pohocco-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell Slope	0.50 0.12
MoD: Monona-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Slope	0.04	Very limited Slope Shrink-swell	1.00 0.50
MoD2: Pohocco-----	100	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Slope	0.04	Very limited Slope Shrink-swell	1.00 0.50
MoE: Monona-----	100	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope	0.96	Very limited Slope Shrink-swell	1.00 0.50
MsE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Pohocco-----	50	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Slope Shrink-swell	1.00 0.50
MsF: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope	1.00	Very limited Slope Shrink-swell	1.00 0.50
MsF2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope	1.00	Very limited Slope Shrink-swell	1.00 0.50

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MsG: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope	1.00	Very limited Slope Shrink-swell	1.00 0.50
On: Onawa-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Pa: Percival-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Pb: Pits-----	100	Not rated		Not rated		Not rated	
Pc: Platte-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
PdD2: Ida-----	50	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Ponca-----	50	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
PdE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
Ra: Riverwash-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Sd: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Sp: Sarpy-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ss: Nodaway Variant-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.35	Very limited Flooding	1.00
StE2: Steinauer-----	100	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope Shrink-swell	1.00 0.50
Wb: Wabash-----	100	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 1.00 1.00
Wc: Wabash-----	100	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 1.00 1.00



BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Wm: Wann-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding	1.00
Wt: Barney-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Shrink-swell	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Too clayey	0.88	Flooding	0.60
		Depth to saturated zone	1.00	Flooding	0.60		
		Frost action	0.50	Cutbanks cave	0.10		
Ac: Albaton-----	100	Very limited Shrink-swell	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Too clayey	0.88	Too clayey	1.00
		Depth to saturated zone	1.00	Flooding	0.60	Flooding	0.60
		Frost action	0.50	Cutbanks cave	0.10		
Af: Alda-----	100	Very limited Frost action	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Flooding	0.60
		Flooding	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.03
		Depth to saturated zone	0.03	Flooding	0.60	Droughty	0.00
Ag: Alda-----	100	Very limited Frost action	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Flooding	0.60
		Flooding	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.03
		Depth to saturated zone	0.03	Flooding	0.60		
Ca: Carr-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.60	Somewhat limited Flooding	0.60
				Cutbanks cave	0.10		
Cc: Cass-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Flooding	0.40				
Cd: Cass-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Flooding	0.40	Depth to saturated zone	0.95		
Ce: Cass-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Flooding	0.40				
Cg: Colo-----	100	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.60	Flooding	0.60
		Depth to saturated zone	1.00	Cutbanks cave	0.10		
		Shrink-swell	0.50				
Ck: Colo-----	50	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.60	Flooding	0.60
		Depth to saturated zone	1.00	Cutbanks cave	0.10		
		Shrink-swell	0.50				
Kennebec-----	50	Very limited Frost action	1.00	Somewhat limited Flooding	0.60	Somewhat limited Flooding	0.60
		Flooding	1.00	Depth to saturated zone	0.35		
		Shrink-swell	0.50	Cutbanks cave	0.10		
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Somewhat limited Slope	0.96	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.96
		Frost action	0.50	Slope	0.96		

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ed: Eudora-----	100	Very limited Frost action Flooding	1.00 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Ga: Gibbon-----	100	Very limited Frost action Flooding Depth to saturated zone	1.00 1.00 0.03	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.60 0.10	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gb: Gibbon-----	100	Very limited Frost action Flooding Depth to saturated zone	1.00 1.00 0.03	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.60 0.10	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gc: Gibbon-----	100	Very limited Frost action Flooding Shrink-swell Depth to saturated zone	1.00 1.00 0.50 0.03	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.60 0.10	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gs: Gibbon-----	65	Very limited Frost action Flooding Shrink-swell Depth to saturated zone	1.00 1.00 0.50 0.03	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.60 0.10	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Saltine-----	35	Very limited Frost action Flooding Shrink-swell Depth to saturated zone	1.00 1.00 0.50 0.03	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.60 0.10	Very limited Sodium content Flooding Depth to saturated zone Salinity	1.00 0.60 0.03 0.00
Gu: Ida Variant-----	100	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Ha: Haynie-----	100	Very limited Frost action	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
IdD2: Ida-----	100	Very limited Frost action Slope	1.00 0.63	Somewhat limited Slope Cutbanks cave	0.63 0.10	Somewhat limited Slope	0.63
IdE: Ida-----	100	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
IdE2: Ida-----	100	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Im: Inavale-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.40
In: Inavale-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.61
JuB: Judson-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ke: Kennebec-----	100	Very limited Frost action Flooding	1.00 1.00	Somewhat limited Flooding Depth to saturated zone Cutbanks cave	0.60 0.35 0.10	Somewhat limited Flooding	0.60
Le: Lockton-----	100	Very limited Frost action Flooding  Depth to saturated zone	1.00 1.00 0.03	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Ls: Luton-----	100	Very limited Shrink-swell  Depth to saturated zone Frost action Flooding	1.00 1.00 0.50 0.40	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.88 0.10	Very limited Depth to saturated zone	1.00
Lt: Luton-----	100	Very limited Shrink-swell  Depth to saturated zone Frost action Flooding	1.00 1.00 0.50 0.40	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.88 0.10	Very limited Depth to saturated zone	1.00
Lu: Luton-----	100	Very limited Shrink-swell  Depth to saturated zone Frost action Flooding	1.00 1.00 0.50 0.40	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.88 0.10	Very limited Depth to saturated zone Too clayey	1.00 1.00
MaA: Marshall-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MaB: Marshall-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MaC: Marshall-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MaC2: Marshall-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MaD: Marshall-----	100	Very limited Frost action Shrink-swell Slope	1.00 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
MeD2: Marshall-----	70	Very limited Frost action Shrink-swell Slope	1.00 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
Ponca-----	30	Very limited Frost action Slope	1.00 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
MfE: Marshall-----	50	Very limited Frost action Slope Shrink-swell	1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96
Ponca-----	50	Very limited Frost action Slope	1.00 0.96	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MfE2: Marshall-----	50	Very limited Frost action Slope Shrink-swell	1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96
Ponca-----	50	Very limited Frost action Slope	1.00 0.96	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96
MoA: Monona-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MoB: Monona-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MoC: Monona-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MoC2: Pohocco-----	100	Very limited Frost action Shrink-swell	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
MoD: Monona-----	100	Very limited Frost action Shrink-swell Slope	1.00 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
MoD2: Pohocco-----	100	Very limited Frost action Shrink-swell Slope	1.00 0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
MoE: Monona-----	100	Very limited Frost action Slope Shrink-swell	1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96
MsE2: Ida-----	50	Very limited Frost action Slope	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Pohocco-----	50	Very limited Frost action Slope Shrink-swell	1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	0.96 0.10	Somewhat limited Slope	0.96
MsF: Ida-----	50	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Monona-----	50	Very limited Slope Frost action Shrink-swell	1.00 1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
MsF2: Ida-----	50	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Monona-----	50	Very limited Slope Frost action Shrink-swell	1.00 1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
MsG: Ida-----	50	Very limited Slope Frost action	1.00 1.00	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Monona-----	50	Very limited Slope Frost action Shrink-swell	1.00 1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
On: Onawa-----	100	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Very limited Too clayey	1.00
		Flooding	1.00	Too clayey	0.88	Flooding	0.60
		Depth to saturated zone	0.03	Flooding	0.60	Depth to saturated zone	0.03
				Cutbanks cave	0.10		
Pa: Percival-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Very limited Too clayey	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Flooding	0.60
		Depth to saturated zone	0.03	Flooding	0.60	Droughty	0.16
				Too clayey	0.50	Depth to saturated zone	0.03
Pb: Pits-----	100	Not rated		Not rated		Not rated	
PC: Platte-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Flooding	0.60
		Frost action	0.50	Depth to saturated zone	1.00	Droughty	0.22
		Depth to saturated zone	0.03	Flooding	0.60	Depth to saturated zone	0.03
PdD2: Ida-----	50	Very limited Frost action	1.00	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16
		Slope	0.16	Cutbanks cave	0.10		
Ponca-----	50	Very limited Frost action	1.00	Somewhat limited Cutbanks cave	0.10	Somewhat limited Slope	0.04
		Slope	0.04	Slope	0.04		
PdE2: Ida-----	50	Very limited Frost action	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	1.00	Cutbanks cave	0.10		
Ponca-----	50	Very limited Frost action	1.00	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
		Slope	0.96	Cutbanks cave	0.10		
Ra: Riverwash-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
				Flooding	0.80	Droughty Too sandy	1.00 0.50
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	1.00	Cutbanks cave	0.10		
Sd: Inavale-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Very limited Flooding	1.00
				Flooding	0.80	Too sandy	0.50
						Droughty	0.34
Sp: Sarpy-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.64
				Flooding	0.60	Flooding	0.60
Ss: Nodaway Variant-----	100	Very limited Frost action	1.00	Somewhat limited Flooding	0.80	Very limited Flooding	1.00
		Flooding	1.00	Depth to saturated zone	0.35		
				Cutbanks cave	0.10		

BUILDING SITE DEVELOPMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
StE2: Steinauer-----	100	Very limited Slope Shrink-swell Frost action	1.00 0.50 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
Wb: Wabash-----	100	Very limited Shrink-swell  Flooding Depth to saturated zone Frost action	1.00  1.00 1.00 0.50	Very limited Depth to saturated zone Flooding Too clayey Cutbanks cave	1.00  0.60 0.50 0.10	Very limited Depth to saturated zone Flooding	1.00  0.60
Wc: Wabash-----	100	Very limited Shrink-swell  Flooding Depth to saturated zone Frost action	1.00  1.00 1.00 0.50	Very limited Depth to saturated zone Flooding Too clayey Cutbanks cave	1.00  0.60 0.50 0.10	Very limited Depth to saturated zone Too clayey Flooding	1.00  1.00 0.60
Wm: Wann-----	100	Very limited Frost action Flooding	1.00 1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding	0.60
Wt: Barney-----	100	Very limited Depth to saturated zone Flooding  Frost action	1.00  1.00  0.50	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00  1.00 0.80	Very limited Flooding  Depth to saturated zone Droughty	1.00  1.00 0.16
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	





CONSTRUCTION MATERIALS  
Douglas and Sarpy Counties, Nebraska

Construction Materials

These tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In the first table, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Ab: Albaton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ac: Albaton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Af: Alda-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.03 0.14
Ag: Alda-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.14
Ca: Carr-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cc: Cass-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.22
Cd: Cass-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.01 0.22
Ce: Cass-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.22
Cg: Colo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ck: Colo-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Kennebec-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cm: Urban Land-----	100	Not rated		Not rated	
DcE: Dickinson-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.04 0.72
Ed: Eudora-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ga: Gibbon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.16
Gb: Gibbon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Gc: Gibbon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Gs: Gibbon-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Saltine-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Gu: Ida Variant-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ha: Haynie-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
IdD2: Ida-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
IdE: Ida-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
IdE2: Ida-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Im: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.33 0.33
In: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.33 0.33
JuB: Judson-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ke: Kennebec-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Le: Lockton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.28
Ls: Luton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lt: Luton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lu: Luton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
MaA: Marshall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MaB: Marshall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MaC: Marshall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MaC2: Marshall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MaD: Marshall-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MeD2: Marshall-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ponca-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MfE: Marshall-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ponca-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MfE2: Marshall-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ponca-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoA: Monona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoB: Monona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoC: Monona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoC2: Pohocco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoD: Monona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
MoD2: Pohocco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MoE: Monona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MsE2: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pohocco-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MsF: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Monona-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MsF2: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Monona-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MsG: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Monona-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
On: Onawa-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pa: Percival-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.19
Pb: Pits-----	100	Not rated		Not rated	
Pc: Platte-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.16
PdD2: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ponca-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
PdE2: Ida-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Ponca-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ra: Riverwash-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.09 0.22
Rk: Rock Outcrop-----	100	Not rated		Not rated	
Rn: Ida Variant-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sd: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.33 0.33
Sp: Sarpy-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.28 0.96
Ss: Nodaway Variant----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
StE2: Steinauer-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Wb: Wabash-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Wc: Wabash-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Wm: Wann-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.07 0.10
Wt: Barney-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.04 0.16
zwa: Water-----	100	Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Poor Too clayey Low content of organic matter Carbonate content Water erosion	0.00 0.50 0.97 0.99	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone Carbonate content	0.00 0.00 0.97
Ac: Albaton-----	100	Poor Too clayey Low content of organic matter Carbonate content	0.00 0.50 0.97	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone Carbonate content	0.00 0.00 0.97
Af: Alda-----	100	Fair Low content of organic matter Droughty Too sandy Sodium content	0.12 0.57 0.62 0.97	Fair Depth to saturated zone	0.76	Fair Rock fragments Too sandy Depth to saturated zone Sodium content	0.50 0.62 0.76 0.98
Ag: Alda-----	100	Fair Low content of organic matter Too sandy Droughty Sodium content	0.12 0.62 0.83 0.97	Fair Depth to saturated zone	0.76	Fair Rock fragments Too sandy Depth to saturated zone Sodium content	0.50 0.62 0.76 0.98
Ca: Carr-----	100	Fair Low content of organic matter	0.12	Good		Good	
Cc: Cass-----	100	Fair Low content of organic matter Too acid	0.88 0.92	Good		Good	
Cd: Cass-----	100	Fair Too acid	0.92	Good		Good	
Ce: Cass-----	100	Fair Low content of organic matter Too acid	0.88 0.92	Good		Good	
Cg: Colo-----	100	Good		Poor Depth to saturated zone Shrink-swell	0.00 0.87	Poor Depth to saturated zone	0.00
Ck: Colo-----	50	Good		Poor Depth to saturated zone Shrink-swell	0.00 0.87	Poor Depth to saturated zone	0.00
Kennebec-----	50	Fair Water erosion	0.90	Fair Shrink-swell	0.87	Good	
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Fair Low content of organic matter Too acid	0.12 0.99	Good		Fair Slope	0.04

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ed: Eudora-----	100	Fair Water erosion	0.90	Good		Good	
Ga: Gibbon-----	100	Poor Wind erosion	0.00	Fair Depth to saturated zone	0.76	Poor Hard to reclaim	0.00
		Too acid	0.54			Depth to saturated zone	0.76
		Low content of organic matter	0.88				
		Water erosion	0.99				
Gb: Gibbon-----	100	Fair Low content of organic matter	0.88	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
Gc: Gibbon-----	100	Fair Low content of organic matter	0.88	Fair Depth to saturated zone Shrink-swell	0.76 0.99	Fair Depth to saturated zone	0.76
Gs: Gibbon-----	65	Good		Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
Saltine-----	35	Poor Sodium content	0.00	Fair Depth to saturated zone	0.76	Poor Sodium content	0.00
		Too alkaline	0.00	Shrink-swell	0.82	Depth to saturated zone	0.76
		Low content of organic matter	0.12				
		Salinity	0.88				
		Water erosion	0.90				
Gu: Ida Variant-----	100	Fair Low content of organic matter	0.12	Poor Slope	0.00	Poor Slope	0.00
		Water erosion	0.90				
		Carbonate content	0.97			Fair Carbonate content	0.97
IdD2: Ida-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.37
		Water erosion	0.90			Carbonate content	0.97
		Carbonate content	0.97				
IdE: Ida-----	100	Fair Low content of organic matter	0.12	Fair Slope	0.02	Poor Slope	0.00
		Water erosion	0.90			Carbonate content	0.97
		Carbonate content	0.97				
IdE2: Ida-----	100	Fair Low content of organic matter	0.12	Fair Slope	0.02	Poor Slope	0.00
		Water erosion	0.90			Carbonate content	0.97
		Carbonate content	0.97				



CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Im: Inavale-----	100	Poor Wind erosion Too sandy Low content of organic matter Droughty	0.00 0.06 0.12 0.13	Good		Fair Too sandy	0.06
In: Inavale-----	100	Poor Wind erosion Too sandy Droughty Low content of organic matter	0.00 0.06 0.06 0.12	Good		Fair Too sandy	0.06
JuB: Judson-----	100	Fair Water erosion	0.90	Fair Shrink-swell	0.87	Good	
Ke: Kennebec-----	100	Fair Water erosion	0.90	Fair Shrink-swell	0.98	Good	
Le: Lockton-----	100	Fair Low content of organic matter	0.12	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
Ls: Luton-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.50 0.90	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
Lt: Luton-----	100	Poor Too clayey Low content of organic matter	0.00 0.50	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
Lu: Luton-----	100	Poor Too clayey Low content of organic matter	0.00 0.50	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
MaA: Marshall-----	100	Fair Low content of organic matter Water erosion Too clayey	0.88 0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey	0.72
MaB: Marshall-----	100	Fair Low content of organic matter Water erosion Too clayey	0.88 0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey	0.72
MaC: Marshall-----	100	Fair Low content of organic matter Water erosion Too clayey	0.88 0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey	0.72
MaC2: Marshall-----	100	Fair Low content of organic matter Water erosion Too clayey	0.12 0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey	0.77

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MaD: Marshall-----	100	Fair Low content of organic matter Water erosion Too clayey	0.12 0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey Slope	0.72 0.96
MeD2: Marshall-----	70	Fair Water erosion Too clayey	0.90 0.99	Fair Shrink-swell	0.87	Fair Too Clayey Slope	0.77 0.96
Ponca-----	30	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Fair Slope	0.96
MfE: Marshall-----	50	Fair Low content of organic matter Water erosion Too clayey	0.12 0.90 0.99	Fair Shrink-swell	0.87	Fair Slope Too Clayey	0.04 0.65
Ponca-----	50	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Fair Slope	0.04
MfE2: Marshall-----	50	Fair Low content of organic matter Water erosion Too clayey	0.12 0.90 0.99	Fair Shrink-swell	0.87	Fair Slope Too Clayey	0.04 0.77
Ponca-----	50	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Fair Slope	0.04
MoA: Monona-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Good	
MoB: Monona-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Good	
MoC: Monona-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Good	
MoC2: Pohocco-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Good	
MoD: Monona-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Fair Slope	0.96
MoD2: Pohocco-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Fair Slope	0.96

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MoE: Monona-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Good		Fair Slope	0.04
MsE2: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Fair Slope	0.98	Poor Slope Carbonate content	0.00 0.97
Pohocco-----	50	Fair Low content of organic matter Water erosion	0.88 0.90	Fair Shrink-swell	0.89	Fair Slope	0.04
MsF: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Poor Slope	0.00	Poor Slope Carbonate content	0.00 0.97
Monona-----	50	Fair Low content of organic matter Water erosion	0.12 0.90	Fair Slope	0.02	Poor Slope	0.00
MsF2: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Poor Slope	0.00	Poor Slope Carbonate content	0.00 0.97
Monona-----	50	Fair Low content of organic matter Water erosion	0.12 0.90	Fair Slope	0.02	Poor Slope	0.00
MsG: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Poor Slope	0.00	Poor Slope Carbonate content	0.00 0.97
Monona-----	50	Fair Low content of organic matter Water erosion	0.12 0.90	Poor Slope	0.00	Poor Slope	0.00
On: Onawa-----	100	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Fair Depth to saturated zone Shrink-swell	0.76 0.99	Fair Depth to saturated zone Carbonate content	0.76 0.97
Pa: Percival-----	100	Fair Low content of organic matter Too sandy Droughty	0.12 0.32 0.74	Fair Depth to saturated zone Shrink-swell	0.76 0.99	Fair Too sandy Depth to saturated zone	0.32 0.76
Pb: Pits-----	100	Not rated		Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PC: Platte-----	100	Fair Low content of organic matter Droughty Too sandy	0.12 0.22 0.50	Fair Depth to saturated zone	0.76	Poor Hard to reclaim Rock fragments Too sandy Depth to saturated zone Hard to reclaim	0.00 0.03 0.50 0.76 0.98
PdD2: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Good		Fair Slope Carbonate content	0.84 0.97
Ponca-----	50	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Fair Slope	0.96
PdE2: Ida-----	50	Fair Low content of organic matter Water erosion Carbonate content	0.12 0.90 0.97	Fair Slope	0.98	Poor Slope Carbonate content	0.00 0.97
Ponca-----	50	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Fair Slope	0.04
Ra: Riverwash-----	100	Poor Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12	Poor Depth to saturated zone	0.00	Poor Hard to reclaim Depth to saturated zone Rock fragments Hard to reclaim	0.00 0.00 0.00 0.32
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	
Rn: Ida Variant-----	100	Fair Low content of organic matter Water erosion	0.12 0.90	Poor Slope	0.00	Poor Slope	0.00
Sd: Inavale-----	100	Poor Wind erosion Too sandy Low content of organic matter Droughty	0.00 0.06 0.12 0.71	Good		Fair Too sandy	0.06
Sp: Sarpy-----	100	Poor Wind erosion Too sandy Droughty Low content of organic matter	0.00 0.11 0.38 0.88	Good		Fair Too sandy	0.11
Ss: Nodaway Variant-----	100	Fair Low content of organic matter Water erosion	0.88 0.90	Good		Good	

CONSTRUCTION MATERIALS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
StE2: Steinauer-----	100	Fair Low content of organic matter Water erosion	0.12 0.99	Fair Slope Shrink-swell	0.32 0.87	Poor Slope	0.00
Wb: Wabash-----	100	Poor Too clayey Too acid	0.00 0.92	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
Wc: Wabash-----	100	Poor Too clayey Too acid	0.00 0.92	Poor Shrink-swell Depth to saturated zone	0.00 0.00	Poor Too Clayey Depth to saturated zone	0.00 0.00
Wm: Wann-----	100	Fair Low content of organic matter	0.88	Fair Depth to saturated zone	0.91	Fair Depth to saturated zone	0.91
Wt: Barney-----	100	Fair Low content of organic matter Droughty Too sandy	0.12 0.37 0.50	Poor Depth to saturated zone	0.00	Poor Hard to reclaim Depth to saturated zone Rock fragments Too sandy Hard to reclaim	0.00 0.00 0.03 0.50 0.98
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	



RECREATIONAL INTERPRETATIONS  
Douglas and Sarpy Counties, Nebraska

## Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00			Flooding	0.60
Ac: Albaton-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Too clayey	1.00	Too clayey	1.00
		Too clayey	1.00			Flooding	0.60
Af: Alda-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Ag: Alda-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Ca: Carr-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Cc: Cass-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Cd: Cass-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Ce: Cass-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Cg: Colo-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Restricted permeability	0.21	Flooding	0.60
		Restricted permeability	0.21			Restricted permeability	0.21
Ck: Colo-----	50	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Restricted permeability	0.21	Flooding	0.60
		Restricted permeability	0.21			Restricted permeability	0.21
Kennebec-----	50	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
Ed: Eudora-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Ga: Gibbon-----	100	Very limited Flooding	1.00	Somewhat limited Too sandy	0.28	Somewhat limited Flooding	0.60
		Too sandy	0.28	Depth to saturated zone	0.03	Too sandy	0.28
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Gb: Gibbon-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07



RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Gc: Gibbon-----	100	Very limited Flooding	1.00	Somewhat limited Restricted permeability	0.15	Somewhat limited Flooding	0.60
		Restricted permeability	0.15	Depth to saturated zone	0.03	Restricted permeability	0.15
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Gs: Gibbon-----	65	Very limited Flooding	1.00	Somewhat limited Restricted permeability	0.15	Somewhat limited Flooding	0.60
		Restricted permeability	0.15	Depth to saturated zone	0.03	Restricted permeability	0.15
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Saltine-----	35	Very limited Flooding	1.00	Very limited Sodium content	1.00	Very limited Sodium content	1.00
		Sodium content	1.00	Restricted permeability	0.21	Flooding	0.60
		Restricted permeability	0.21	Depth to saturated zone	0.03	Restricted permeability	0.21
		Depth to saturated zone	0.07	Salinity	0.00	Depth to saturated zone	0.07
		Salinity	0.00			Salinity	0.00
Gu: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ha: Haynie-----	100	Not limited		Not limited		Not limited	
IdD2: Ida-----	100	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
IdE: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
IdE2: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Im: Inavale-----	100	Somewhat limited Too sandy	0.50	Somewhat limited Too sandy	0.50	Somewhat limited Too sandy	0.50
In: Inavale-----	100	Somewhat limited Too sandy	0.50	Somewhat limited Too sandy	0.50	Somewhat limited Slope Too sandy	0.87 0.50
JuB: Judson-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Slope	0.87
						Restricted permeability	0.21
Ke: Kennebec-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Le: Lockton-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Ls: Luton-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00				
Lt: Luton-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00				
Lu: Luton-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Restricted permeability	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Too clayey	1.00	Too clayey	1.00
		Too clayey	1.00				
MaA: Marshall-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21
MaB: Marshall-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability Slope	0.21 0.00
MaC: Marshall-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Slope	0.87
						Restricted permeability	0.21
MaC2: Marshall-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Slope	0.87
						Restricted permeability	0.21
MaD: Marshall-----	100	Somewhat limited Restricted permeability Slope	0.21 0.04	Somewhat limited Restricted permeability Slope	0.21 0.04	Very limited Slope	1.00
						Restricted permeability	0.21
MeD2: Marshall-----	70	Somewhat limited Restricted permeability Slope	0.21 0.04	Somewhat limited Restricted permeability Slope	0.21 0.04	Very limited Slope	1.00
						Restricted permeability	0.21
Ponca-----	30	Somewhat limited Restricted permeability Slope	0.21 0.04	Somewhat limited Restricted permeability Slope	0.21 0.04	Very limited Slope	1.00
						Restricted permeability	0.21
MfE: Marshall-----	50	Somewhat limited Slope Restricted permeability	0.96 0.21	Somewhat limited Slope Restricted permeability	0.96 0.21	Very limited Slope Restricted permeability	1.00 0.21
Ponca-----	50	Somewhat limited Slope Restricted permeability	0.96 0.21	Somewhat limited Slope Restricted permeability	0.96 0.21	Very limited Slope Restricted permeability	1.00 0.21
MfE2: Marshall-----	50	Somewhat limited Slope Restricted permeability	0.96 0.21	Somewhat limited Slope Restricted permeability	0.96 0.21	Very limited Slope Restricted permeability	1.00 0.21
Ponca-----	50	Somewhat limited Slope Restricted permeability	0.96 0.21	Somewhat limited Slope Restricted permeability	0.96 0.21	Very limited Slope Restricted permeability	1.00 0.21
MoA: Monona-----	100	Not limited		Not limited		Not limited	
MoB: Monona-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
MoC: Monona-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
MoC2: Pohocco-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Somewhat limited Slope	0.87
						Restricted permeability	0.21
MoD: Monona-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
MoD2: Pohocco-----	100	Somewhat limited Restricted permeability	0.21	Somewhat limited Restricted permeability	0.21	Very limited Slope	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Slope	0.04	Slope	0.04	Restricted permeability	0.21
MoE: Monona-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
MsE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Pohocco-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
		Restricted permeability	0.21	Restricted permeability	0.21	Restricted permeability	0.21
MsF: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
MsF2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
MsG: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
On: Onawa-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability Too clayey	1.00	Too clayey	1.00	Too clayey	1.00
		Depth to saturated zone	0.07	Depth to saturated zone	0.03	Flooding	0.60
						Depth to saturated zone	0.07
Pa: Percival-----	100	Very limited Flooding	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Restricted permeability Too clayey	1.00	Too clayey	1.00	Too clayey	1.00
		Depth to saturated zone	0.07	Depth to saturated zone	0.03	Flooding	0.60
						Depth to saturated zone	0.07
Pb: Pits-----	100	Not rated		Not rated		Not rated	
PC: Platte-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
PdD2: Ida-----	50	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Ponca-----	50	Somewhat limited Restricted permeability Slope	0.21	Somewhat limited Restricted permeability Slope	0.21	Very limited Slope	1.00
			0.04		0.04	Restricted permeability	0.21
PdE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
		Restricted permeability	0.21	Restricted permeability	0.21	Restricted permeability	0.21
Ra: Riverwash-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Too sandy	0.50	Flooding	1.00
		Too sandy	0.50	Flooding	0.40	Too sandy	0.50
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Sd: Inavale-----	100	Very limited Flooding Too sandy	1.00 0.72	Somewhat limited Too sandy Flooding	0.72 0.40	Very limited Flooding Too sandy	1.00 0.72
Sp: Sarpy-----	100	Very limited Flooding Too sandy	1.00 1.00	Very limited Too sandy	1.00	Very limited Too sandy Flooding	1.00 0.60
Ss: Nodaway Variant----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
StE2: Steinauer-----	100	Very limited Slope Restricted permeability	1.00 0.15	Very limited Slope Restricted permeability	1.00 0.15	Very limited Slope Restricted permeability	1.00 0.15
Wb: Wabash-----	100	Very limited Flooding  Restricted permeability Depth to saturated zone	1.00  1.00 1.00	Very limited Restricted permeability Depth to saturated zone	1.00  1.00	Very limited Restricted permeability Depth to saturated zone Flooding	1.00  1.00 0.60
Wc: Wabash-----	100	Very limited Flooding  Restricted permeability Depth to saturated zone Too clayey	1.00  1.00 1.00 1.00	Very limited Restricted permeability Depth to saturated zone Too clayey	1.00  1.00 1.00	Very limited Restricted permeability Depth to saturated zone Too clayey Flooding	1.00  1.00 1.00 0.60
Wm: Wann-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Wt: Barney-----	100	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60
Ac: Albaton-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00	Very limited Depth to saturated zone Too clayey Flooding	1.00 1.00 0.60
Af: Alda-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone Droughty	0.60 0.03 0.00
Ag: Alda-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Ca: Carr-----	100	Not limited		Somewhat limited Flooding	0.60
Cc: Cass-----	100	Not limited		Not limited	
Cd: Cass-----	100	Not limited		Not limited	
Ce: Cass-----	100	Not limited		Not limited	
Cg: Colo-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60
Ck: Colo-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60
Kennebec-----	50	Not limited		Somewhat limited Flooding	0.60
Cm: Urban Land-----	100	Not rated		Not rated	
DcE: Dickinson-----	100	Not limited		Somewhat limited Slope	0.96
Ed: Eudora-----	100	Not limited		Not limited	
Ga: Gibbon-----	100	Somewhat limited Too sandy	0.28	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gb: Gibbon-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gc: Gibbon-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Gs: Gibbon-----	65	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Saltine-----	35	Not limited		Very limited Sodium content Flooding Depth to saturated zone Salinity	1.00 0.60 0.03 0.00
Gu: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ha: Haynie-----	100	Not limited		Not limited	
IdD2: Ida-----	100	Not limited		Somewhat limited Slope	0.63
IdE: Ida-----	100	Somewhat limited Slope	0.98	Very limited Slope	1.00
IdE2: Ida-----	100	Somewhat limited Slope	0.98	Very limited Slope	1.00
Im: Inavale-----	100	Somewhat limited Too sandy	0.50	Somewhat limited Droughty	0.40
In: Inavale-----	100	Somewhat limited Too sandy	0.50	Somewhat limited Droughty	0.61
JuB: Judson-----	100	Not limited		Not limited	
Ke: Kennebec-----	100	Not limited		Somewhat limited Flooding	0.60
Le: Lockton-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Ls: Luton-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Lt: Luton-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Lu: Luton-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00	Very limited Depth to saturated zone Too clayey	1.00 1.00
MaA: Marshall-----	100	Not limited		Not limited	
MaB: Marshall-----	100	Not limited		Not limited	
MaC: Marshall-----	100	Not limited		Not limited	
MaC2: Marshall-----	100	Not limited		Not limited	
MaD: Marshall-----	100	Not limited		Somewhat limited Slope	0.04
MeD2: Marshall-----	70	Not limited		Somewhat limited Slope	0.04
Ponca-----	30	Not limited		Somewhat limited Slope	0.04
MfE: Marshall-----	50	Not limited		Somewhat limited Slope	0.96
Ponca-----	50	Not limited		Somewhat limited Slope	0.96
MfE2: Marshall-----	50	Not limited		Somewhat limited Slope	0.96
Ponca-----	50	Not limited		Somewhat limited Slope	0.96
MoA: Monona-----	100	Not limited		Not limited	
MoB: Monona-----	100	Not limited		Not limited	
MoC: Monona-----	100	Not limited		Not limited	
MoC2: Pohocco-----	100	Not limited		Not limited	
MoD: Monona-----	100	Not limited		Somewhat limited Slope	0.04
MoD2: Pohocco-----	100	Not limited		Somewhat limited Slope	0.04

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
MoE: Monona-----	100	Not limited		Somewhat limited Slope	0.96
MsE2: Ida-----	50	Somewhat limited Slope	0.02	Very limited Slope	1.00
Pohocco-----	50	Not limited		Somewhat limited Slope	0.96
MsF: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Somewhat limited Slope	0.98	Very limited Slope	1.00
MsF2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Somewhat limited Slope	0.98	Very limited Slope	1.00
MsG: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00
On: Onawa-----	100	Very limited Too clayey	1.00	Very limited Too clayey Flooding Depth to saturated zone	1.00 0.60 0.03
Pa: Percival-----	100	Very limited Too clayey	1.00	Very limited Too clayey Flooding Droughty Depth to saturated zone	1.00 0.60 0.16 0.03
Pb: Pits-----	100	Not rated		Not rated	
Pc: Platte-----	100	Not limited		Somewhat limited Flooding Droughty Depth to saturated zone	0.60 0.22 0.03
PdD2: Ida-----	50	Not limited		Somewhat limited Slope	0.16
Ponca-----	50	Not limited		Somewhat limited Slope	0.04
PdE2: Ida-----	50	Somewhat limited Slope	0.02	Very limited Slope	1.00
Ponca-----	50	Not limited		Somewhat limited Slope	0.96
Ra: Riverwash-----	100	Very limited Depth to saturated zone Too sandy Flooding	1.00 0.50 0.40	Very limited Flooding Depth to saturated zone Droughty Too sandy	1.00 1.00 1.00 0.50
Rk: Rock Outcrop-----	100	Not rated		Not rated	
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00
Sd: Inavale-----	100	Somewhat limited Too sandy Flooding	0.72 0.40	Very limited Flooding Too sandy Droughty	1.00 0.50 0.34
Sp: Sarpy-----	100	Very limited		Somewhat limited	

RECREATIONAL INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
		Too sandy	1.00	Droughty Flooding	0.64 0.60
Ss: Nodaway Variant-----	100	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
StE2: Steinauer-----	100	Somewhat limited Slope	0.68	Very limited Slope	1.00
Wb: Wabash-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60
Wc: Wabash-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00	Very limited Depth to saturated zone Too clayey Flooding	1.00 1.00 0.60
Wm: Wann-----	100	Not limited		Somewhat limited Flooding	0.60
Wt: Barney-----	100	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 0.16
zwa: Water-----	100	Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated	





WILDLIFE INTERPRETATIONS  
Douglas and Sarpy Counties, Nebraska

Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, brome grass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS  
Douglas and Sarpy Counties, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Ab: ALBATON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Ac: ALBATON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Af: ALDA-----	Fair	Fair	Fair	Good	Good	Good	Fair	Fair	Fair	Good	Fair	Good
Ag: ALDA-----	Fair	Fair	Fair	Good	Good	Good	Fair	Fair	Fair	Good	Fair	Good
Ca: CARR-----	Good	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor	---
Cc: CASS-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
Cd: CASS-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
Ce: CASS-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
Cg: COLO-----	Good	Fair	Good	Fair	Poor	---	Good	Good	Fair	Fair	Good	---
Ck: COLO-----	Good	Fair	Good	Fair	Poor	---	Good	Good	Fair	Fair	Good	---
KENNEBEC-----	Good	Good	Good	Good	Good	---	Poor	Poor	Good	Good	Poor	---
Cm: URBAN LAND-----	---	---	---	---	---	---	---	---	---	---	---	---
DcE: DICKINSON-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
Ed: EUDORA-----	Good	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor	---
Ga: GIBBON-----	Poor	Fair	Good	Good	Fair	Good	Fair	Good	Fair	Fair	Fair	Good
Gb: GIBBON-----	Good	Good	Good	Good	Fair	Good	Fair	Good	Good	Good	Fair	Good
Gc: GIBBON-----	Good	Good	Good	Good	Fair	Good	Fair	Good	Good	Good	Fair	Good
Gs: GIBBON-----	Good	Good	Good	Good	Fair	Good	Fair	Good	Good	Good	Fair	Good
SALTINE-----	Poor	Poor	Good	Poor	Poor	Poor	Good	Good	Poor	Poor	Good	Poor
Gu: IDA VARIANT-----	Very poor	Very poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
Ha: HAYNIE-----	Good	Good	Good	Good	Good	---	Poor	Poor	Good	Good	Poor	---
IdD2: IDA-----	Fair	Good	Good	Fair	Fair	---	Very poor	Very poor	Good	Fair	Very poor	---
IdE: IDA-----	Poor	Fair	Good	Poor	Poor	---	Very poor	Very poor	Fair	Fair	Very poor	---
IdE2: IDA-----	Poor	Fair	Good	Poor	Poor	---	Very poor	Very poor	Fair	Fair	Very poor	---
Im: INVALE-----	Fair	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Good

WILDLIFE INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
In: INVALE-----	Poor	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
JuB: JUDSON-----	Good	Good	Good	Good	Good	---	Poor	Poor	Good	Good	Poor	---
Ke: KENNEBEC-----	Good	Good	Good	Good	Good	---	Poor	Poor	Good	Good	Poor	---
Le: LOCKTON-----	Fair	Fair	Good	Good	Good	Good	Poor	Poor	Fair	Good	Poor	Good
Ls: LUTON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Lt: LUTON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Lu: LUTON-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
MaA: MARSHALL-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MaB: MARSHALL-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MaC: MARSHALL-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MaC2: MARSHALL-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MaD: MARSHALL-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MeD2: MARSHALL-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
PONCA-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
MfE: MARSHALL-----	Poor	Fair	Good	Good	Good	---	Very poor	Very poor	Fair	Good	Very poor	---
PONCA-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
MfE2: MARSHALL-----	Poor	Fair	Good	Good	Good	---	Very poor	Very poor	Fair	Good	Very poor	---
PONCA-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
MoA: MONONA-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MoB: MONONA-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MoC: MONONA-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MoC2: POHOCCO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good

WILDLIFE INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
MoD: MONONA-----	Good	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MoD2: POHOCCO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
MoE: MONONA-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
MsE2: IDA-----	Fair	Good	Good	Fair	Fair	---	Very poor	Very poor	Good	Fair	Very poor	---
POHOCCO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
MsF: IDA-----	Poor	Fair	Good	Poor	Poor	---	Very poor	Very poor	Fair	Fair	Very poor	---
MONONA-----	Poor	Fair	Good	Fair	Fair	---	Very poor	Very poor	Fair	Fair	Very poor	---
MsF2: IDA-----	Poor	Fair	Good	Poor	Poor	---	Very poor	Very poor	Fair	Fair	Very poor	---
MONONA-----	Poor	Fair	Good	Fair	Fair	---	Very poor	Very poor	Fair	Fair	Very poor	---
MsG: IDA-----	Very poor	Very poor	Good	Poor	Poor	---	Very poor	Very poor	Poor	Poor	Very poor	---
MONONA-----	Very poor	Poor	Good	Fair	Fair	---	Very poor	Very poor	Fair	Fair	Very poor	---
On: ONAWA-----	Fair	Fair	Fair	Poor	Very poor	---	Good	Good	Fair	Poor	Good	---
Pa: PERCIVAL-----	Fair	Fair	Fair	Fair	Poor	---	Fair	Fair	Fair	Fair	Fair	---
Pb: PITS-----	Very poor	Very poor	Poor	Poor	Poor	Poor	Very poor	Fair	Very poor	Very poor	Poor	Poor
Pc: PLATTE-----	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Good	Fair	Good	Fair	Fair
PdD2: IDA-----	Fair	Good	Good	Good	Good	---	Very poor	Very poor	Good	Good	Very poor	---
PONCA-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
PdE2: IDA-----	Fair	Good	Good	Fair	Fair	---	Very poor	Very poor	Good	Fair	Very poor	---
PONCA-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
Ra: RIVERWASH-----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Poor	Very poor	Very poor	Poor	Very poor
Rk: ROCK OUTCROP----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
Rn: IDA VARIANT-----	Very poor	Very poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
Sd: INVALE-----	Very poor	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued  
Douglas and Sarpy Counties, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Sp: SARPY-----	Poor	Poor	Fair	Poor	Poor	---	Very poor	Very poor	Poor	Poor	Very poor	---
Ss: NODAWAY VARIANT-	Poor	Fair	Fair	Fair	Poor	---	Fair	Fair	Fair	Fair	Poor	---
StE2: STEINAUER-----	Poor	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good
Wb: WABASH-----	Poor	Poor	Poor	Poor	Poor	---	Good	Good	Poor	Poor	Good	---
Wc: WABASH-----	Poor	Poor	Poor	Poor	Poor	---	Poor	Good	Poor	Poor	Fair	---
Wm: WANN-----	Good	Good	Good	Good	Fair	Good	Poor	Fair	Good	Good	Fair	Good
Wt: BARNEY-----	Very poor	Poor	Fair	Poor	Poor	Poor	Good	Good	Poor	Fair	Good	Fair
zwa: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---
zwb: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---





YIELDS PER ACRE OF PASTURE AND HAYLAND  
Douglas and Sarpy Counties, Nebraska

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. One animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 2 OF 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Ab: Albaton-----	3w	---	5.00	---
Ac: Albaton-----	3w	3w	4.80	---
Af: Alda-----	3w	3w	2.50	---
Ag: Alda-----	3w	3w	4.00	---
Ca: Carr-----	2w	2w	5.00	---
Cc: Cass-----	2e	2e	4.80	---
Cd: Cass-----	2e	2e	5.00	---
Ce: Cass-----	1	1	5.00	---
Cg: Colo-----	2w	2w	5.50	---
Ck: Colo-----	2w	2w	5.50	---
Kennebec-----	2w	---	5.50	---
Cm: Urban Land-----	---	---	---	---
DcE: Dickinson-----	6e	---	2.00	---
Ed: Eudora-----	1	1	6.00	---
Ga: Gibbon-----	3w	3w	3.00	---
Gb: Gibbon-----	2w	2w	5.00	---
Gc: Gibbon-----	2w	2w	4.50	---
Gs: Gibbon-----	2w	2w	---	---
Saltine-----	6s	---	---	---
Gu: Ida Variant-----	7e	---	2.00	---
Ha: Haynie-----	2w	2w	5.50	---
IdD2: Ida-----	4e	---	3.50	---
IdE: Ida-----	6e	---	---	---
IdE2: Ida-----	6e	---	2.80	---
Im: Inavale-----	4e	3e	---	---
In: Inavale-----	6e	4e	2.20	---
JuB: Judson-----	2e	3e	5.00	---

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 3 OF 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Ke: Kennebec-----	2w	2w	5.50	---
Le: Lockton-----	3w	3w	4.50	---
Ls: Luton-----	3w	3w	4.80	---
Lt: Luton-----	3w	3w	4.50	---
Lu: Luton-----	3w	3w	4.50	---
MaA: Marshall-----	1	1	5.00	---
MaB: Marshall-----	2e	2e	5.00	---
MaC: Marshall-----	2e	3e	4.80	---
MaC2: Marshall-----	3e	2e	4.50	---
MaD: Marshall-----	3e	4e	4.50	---
MeD2: Marshall-----	3e	4e	4.20	---
Ponca-----	3e	4e	4.20	---
MfE: Marshall-----	4e	---	4.00	---
Ponca-----	4e	---	4.00	---
MfE2: Marshall-----	4e	---	4.00	---
Ponca-----	4e	---	4.00	---
MoA: Monona-----	1	1	5.00	---
MoB: Monona-----	2e	2e	5.00	---
MoC: Monona-----	2e	3e	5.00	---
MoC2: Pohocco-----	2e	3e	4.80	---
MoD: Monona-----	3e	4e	4.50	---
MoD2: Pohocco-----	3e	4e	4.20	---
MoE: Monona-----	4e	---	4.00	---
MsE2: Ida-----	4e	---	3.00	---
Pohocco-----	4e	---	3.00	---
MsF: Ida-----	6e	---	2.80	---
Monona-----	6e	---	2.80	---
MsF2: Ida-----	6e	---	---	---
Monona-----	6e	---	---	---

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 4 OF 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
MsG:				
Ida-----	7e	---	2.00	---
Monona-----	7e	---	2.00	---
On:				
Onawa-----	2w	2w	5.00	---
Pa:				
Percival-----	2w	2w	3.50	---
Pb:				
Pits-----	8s	---	---	---
Pc:				
Platte-----	4w	4w	---	---
PdD2:				
Ida-----	3e	4e	4.50	---
Ponca-----	3e	4e	4.50	---
PdE2:				
Ida-----	4e	---	3.00	---
Ponca-----	4e	---	3.00	---
Ra:				
Riverwash-----	8w	---	---	---
Rk:				
Rock Outcrop-----	8s	---	---	---
Rn:				
Ida Variant-----	7e	---	2.00	---
Sd:				
Inavale-----	6w	---	1.80	---
Sp:				
Sarpy-----	6s	4s	---	---
Ss:				
Nodaway Variant-----	6w	---	---	---
StE2:				
Steinauer-----	6e	---	2.70	---
Wb:				
Wabash-----	3w	3w	4.20	---
Wc:				
Wabash-----	3w	3w	4.00	---
Wm:				
Wann-----	2w	2w	3.50	---
Wt:				
Barney-----	5w	---	---	---
zwa:				
Water-----	---	---	---	---
zwb:				
Water-----	---	---	---	---



CONSERVATION TREE AND SHRUB MANAGEMENT  
Douglas and Sarpy Counties,  
Nebraska

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsuitable indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsuited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT  
Douglas and Sarpy  
Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Ab: Albaton-----		Well suited	Well suited	Well suited	Well suited	High Wetness Lime Soil reaction
Ac: Albaton-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness Lime Soil reaction
Af: Alda-----		Well suited	Well suited	Well suited	Well suited	Low
Ag: Alda-----		Well suited	Well suited	Well suited	Well suited	Low
Ca: Carr-----		Well suited	Well suited	Well suited	Well suited	Low
Cc: Cass-----		Well suited	Well suited	Well suited	Well suited	Low
Cd: Cass-----		Well suited	Well suited	Well suited	Well suited	Low
Ce: Cass-----		Well suited	Well suited	Well suited	Well suited	Low
Cg: Colo-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness
Ck: Colo-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness Low
Cm: Urban Land-----		Not rated	Not rated	Not rated	Not rated	Not rated
DcE: Dickinson-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Ed: Eudora-----		Well suited	Well suited	Well suited	Well suited	Low
Ga: Gibbon-----		Well suited	Well suited	Well suited	Well suited	Low
Gb: Gibbon-----		Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Gc: Gibbon-----		Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Gs: Gibbon-----		Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Saltine-----		Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction High Soil reaction Salinity
Gu: Ida Variant-----		Moderately suited Slope	Unsuited Slope	Unsuited Slope	Unsuited Slope	Low
Ha: Haynie-----		Well suited	Well suited	Well suited	Well suited	Moderate Lime
IdD2: Ida-----		Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Lime
IdE: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
IdE2: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Im: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
In: Inavale-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Douglas and Sarpy  
Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
JuB: Judson-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Ke: Kennebec-----		Well suited	Well suited	Well suited	Well suited	Low
Le: Lockton-----		Well suited	Well suited	Well suited	Well suited	Low
LS: Luton-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
Lt: Luton-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness
Lu: Luton-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
MaA: Marshall-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
MaB: Marshall-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
MaC: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
MaC2: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
MaD: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
MeD2: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
Ponca-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MfE: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
Ponca-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MfE2: Marshall-----		Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
Ponca-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MoA: Monona-----		Well suited	Well suited	Well suited	Well suited	Low
MoB: Monona-----		Well suited	Well suited	Well suited	Well suited	Low
MoC: Monona-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MoC2: Pohocco-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MoD: Monona-----		Well suited	Moderately suited	Well suited	Well suited	Low



CONSERVATION TREE AND SHRUB MANAGEMENT  
Douglas and Sarpy  
Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
MoD2: Pohocco-----		Well suited	Slope Moderately suited Slope	Well suited	Well suited	Low
MoE: Monona-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MsE2: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Pohocco-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
MsF: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Monona-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low
MsF2: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Monona-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low
MsG: Ida-----		Moderately suited Slope	Unsuited Slope	Unsuited Slope	Unsuited Slope	Moderate Lime
Monona-----		Moderately suited Slope	Unsuited Slope	Poorly suited Slope	Poorly suited Slope	Low
On: Onawa-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Moderate Lime
Pa: Percival-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Pb: Pits-----		Not rated	Not rated	Not rated	Not rated	Not rated
Pc: Platte-----		Well suited	Well suited	Well suited	Well suited	Low
PdD2: Ida-----		Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Lime
Ponca-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
PdE2: Ida-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Ponca-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Ra: Riverwash-----		Moderately suited Sandiness	Moderately suited Rock fragments Sandiness	Well suited	Well suited	High Wetness Soil reaction
Rk: Rock Outcrop-----		Not rated	Not rated	Not rated	Not rated	Not rated
Rn: Ida Variant-----		Moderately suited Slope	Unsuited Slope	Unsuited Slope	Unsuited Slope	Low
Sd: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
Sp: Sarpy-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Moderate Wetness
Ss: Nodaway Variant-----		Well suited	Well suited	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Douglas and Sarpy  
Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
StE2: Steinauer-----		Moderately suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Poorly suited Slope	Moderate Soil reaction
Wb: Wabash-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Wc: Wabash-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
Wm: Wann-----		Well suited	Well suited	Well suited	Well suited	Low
Wt: Barney-----		Well suited	Well suited	Well suited	Well suited	High Wetness
zwa: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated
zwb: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated



ENGINEERING INDEX PROPERTIES  
Douglas and Sarpy Counties,  
Nebraska

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued  
Douglas and Sarpy Counties,  
Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
Ab: Albaton-----	0-18	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-15
	18-60	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
Ac: Albaton-----	0-6	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
	6-60	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
Af: Alda-----	0-13	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	90-100	85-100	70-100	30-50	15-20	NP-5
	13-20	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	95-100	95-100	70-100	30-50	15-26	NP-7
	20-60	Coarse sand	SC-SM, SM, SP, SP-SM	A-1, A-2, A-3	0	0	70-100	65-95	30-95	2-15	15-20	NP-5
Ag: Alda-----	0-13	Very fine sandy loam	CL, CL-ML, ML	A-4	0	0	90-100	85-100	85-100	50-75	15-25	3-10
	13-20	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	95-100	95-100	70-100	30-50	15-26	NP-7
	20-60	Coarse sand	SC-SM, SM, SP, SP-SM	A-1, A-2, A-3	0	0	70-100	65-95	30-95	2-15	15-20	NP-5
Ca: Carr-----	0-15	Fine sandy loam	SC, SM, CL, ML	A-2, A-4	0	0	100	95-100	70-100	25-75	10-25	2-10
	15-60	Stratified fine sandy loam to loam	CL, ML, SC, SM	A-4	0	0	100	95-100	70-100	35-65	10-25	2-10
Cc: Cass-----	0-16	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-40	15-20	NP-5
	16-39	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-50	15-20	NP-5
	39-60	Coarse sand	SM, SP-SM	A-2, A-3	0	0	95-100	95-100	50-75	5-30	5-15	NP-5
Cd: Cass-----	0-20	Fine sandy loam	SC-SM, SM	A-4	0	0	95-100	95-100	70-85	40-50	0-20	NP-5
	20-30	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	95-100	95-100	85-95	20-40	0-20	NP-5
	30-48	Silty clay loam	CL	A-6	0	0	100	100	95-100	85-95	25-40	11-20
	48-60	Coarse sand	SM, SP-SM	A-2, A-3	0	0	95-100	95-100	50-75	5-30	10-15	NP-5
Ce: Cass-----	0-16	Very fine sandy loam	CL, CL-ML, ML	A-4	0	0	100	100	85-95	50-65	22-35	3-10
	16-39	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-50	15-20	NP-5
	39-60	Coarse sand	SM, SP-SM	A-2, A-3	0	0	95-100	95-100	50-75	5-30	5-15	NP-5
Cg: Colo-----	0-23	Silty clay loam	CH, CL	A-7	0	0	100	100	90-100	90-100	40-60	15-30
	23-40	Silty clay loam	CH, CL	A-7	0	0	100	100	90-100	90-100	40-55	20-30
	40-60	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	80-100	40-55	15-30
Ck: Colo-----	0-23	Silty clay loam	CH, CL	A-7	0	0	100	100	90-100	90-100	40-60	15-30
	23-40	Silty clay loam	CH, CL	A-7	0	0	100	100	90-100	90-100	40-55	20-30
	40-60	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	80-100	40-55	15-30
Kennebec-----	0-17	Silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	25-45	10-20
	17-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-15
Cm: Urban Land----	0-60	Variable			---	---	---	---	---	---	0-14	---
DcE: Dickinson-----	0-10	Fine sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	100	85-95	30-50	15-30	NP-10
	10-16	Fine sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	100	85-95	30-50	15-30	NP-10
	16-30	Fine sandy loam	SC, SC-SM, SM	A-4	0	0	100	100	85-95	35-50	15-30	NP-10
	30-60	Loamy sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	80-95	5-20	10-20	NP-5
Ed: Eudora-----	0-13	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	70-90	20-30	5-10
	13-60	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	75-100	45-90	15-30	NP-10
Ga: Gibbon-----	0-17	Loamy sand	SC-SM, SM	A-2	0	0	100	100	50-75	15-30	5-20	NP-7
	17-60	Loam	CL	A-4, A-6	0	0	100	100	85-95	60-75	25-45	5-20
Gb: Gibbon-----	0-16	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	85-100	70-90	20-30	2-10
	16-40	Silt loam	CL	A-6	0	0	100	100	90-100	80-90	25-38	12-20
	40-60	Stratified very fine sandy loam to silt loam	CL, ML, SC, SM	A-4	0	0	100	100	70-95	35-90	15-25	NP-8
Gc: Gibbon-----	0-16	Silty clay loam	CL	A-6	0	0	100	100	95-100	85-100	32-40	10-15
	16-40	Silty clay loam	CL	A-7	0	0	100	100	95-100	85-95	40-45	20-25
	40-60	Stratified very fine sandy loam to silty clay loam	CL-ML, ML, SC, SM	A-4, A-6	0	0	100	100	75-95	35-95	15-35	NP-15
Gs: Gibbon-----	0-20	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	85-100	70-90	20-30	2-10
	20-40	Silty clay loam	CL	A-7	0	0	100	100	95-100	85-95	40-45	20-25
	40-60	Stratified very fine sandy loam to silty clay loam	CL-ML, ML, SC, SM	A-4, A-6	0	0	100	100	75-95	35-95	15-35	NP-15
Saltine-----	0-6	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	95-100	95-100	85-100	60-90	20-35	3-15
	6-15	Silty clay loam	CL, CL-ML, ML	A-4, A-6, A-7	0	0	95-100	95-100	85-100	60-100	25-55	5-25
	15-20	Silty clay loam	CL, CL-ML, CH	A-4, A-6, A-7	0	0	95-100	95-100	95-100	70-95	25-65	5-35
	20-60	Silty clay loam	CL	A-4, A-6, A-7	0	0	100	100	95-100	70-95	25-50	7-25

ENGINEERING INDEX PROPERTIES--Continued  
Douglas and Sarp County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
Gu:	In											
Ida Variant----	0-6	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	6-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
Ha:												
Haynie-----	0-9	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	85-100	70-100	25-40	5-15
	9-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	85-100	85-100	25-35	5-15
IdD2:												
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
IdE:												
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
IdE2:												
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
Im:												
Inavale-----	0-11	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	85-95	5-35	15-25	NP-5
	11-28	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	28-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
In:												
Inavale-----	0-8	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	85-95	5-35	15-25	NP-5
	8-25	Loamy fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	25-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
JuB:												
Judson-----	0-26	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-35	5-15
	26-37	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-50	15-25
	37-60	Silty clay loam	CL, CL-ML	A-4, A-6, A-7	0	0	100	100	100	95-100	25-50	5-25
Ke:												
Kennebec-----	0-26	Silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	25-45	10-20
	26-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-15
Le:												
Lockton-----	0-21	Silt loam	CL, CL-ML, ML	A-6	0	0	95-100	95-100	90-100	70-95	20-40	3-20
	21-28	Sandy loam	CL, CL-ML, ML	A-4, A-6	0	0	95-100	95-100	85-100	60-95	25-45	5-18
	28-60	Gravelly sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	85-95	65-95	30-65	3-14	15-20	NP-5
Ls:												
Luton-----	0-11	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-15
	11-27	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	27-60	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
Lt:												
Luton-----	0-13	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	95-100	40-60	15-30
	13-27	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	27-60	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
Lu:												
Luton-----	0-17	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	17-27	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	27-60	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
MaA:												
Marshall-----	0-16	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	16-39	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	39-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
MaB:												
Marshall-----	0-14	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	14-38	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	38-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
MaC:												
Marshall-----	0-12	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	12-37	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	37-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
MaC2:												
Marshall-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	8-32	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	32-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
MaD:												
Marshall-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	8-32	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	32-60	Silt loam	CL	A-7, A-6	0	0	100	100	100	95-100	35-50	15-25
MeD2:												
Marshall-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	8-36	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	36-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
Ponca-----	0-5	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	5-20	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	20-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
MfE:												
Marshall-----	0-12	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	12-35	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	35-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
Ponca-----	0-7	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	7-22	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	22-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25

ENGINEERING INDEX PROPERTIES--Continued  
Douglas and Sarpy Counties,  
Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
MfE2: Marshall-----	0-7	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	7-31	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	31-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
Ponca-----	0-4	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	4-20	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	20-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
MoA: Monona-----	0-12	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	12-35	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	35-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
MoB: Monona-----	0-10	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	10-33	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	33-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
MoC: Monona-----	0-10	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	10-33	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	33-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
MoC2: Pohocco-----	0-7	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
	7-30	Silty clay loam	CL, ML	A-6, A-7	0	0	100	100	100	95-100	35-50	10-25
	30-60	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
MoD: Monona-----	0-10	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	10-33	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	33-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
MoD2: Pohocco-----	0-6	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
	6-29	Silty clay loam	CL, ML	A-6, A-7	0	0	100	100	100	95-100	35-50	10-25
	29-60	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
MoE: Monona-----	0-10	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	10-33	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	33-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
MsE2: Pohocco-----	0-6	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
	6-29	Silty clay loam	CL, ML	A-6, A-7	0	0	100	100	100	95-100	35-50	10-25
	29-30	Silt loam	ML	A-4, A-6, A-7	0	0	100	100	100	95-100	30-45	5-15
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
MsF: Monona-----	0-8	Silt loam	CL, ML	A-7, A-6	0	0	100	100	95-100	95-100	35-50	10-25
	8-31	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	31-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
MsF2: Monona-----	0-5	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	5-27	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	27-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
MsG: Monona-----	0-7	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	7-30	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-50	10-25
	30-60	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
On: Onawa-----	0-7	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
	7-18	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	40-60
	18-60	Very fine sandy loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	85-100	25-40	5-20
Pa: Percival-----	0-7	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	7-16	Silty clay	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-60
	16-60	Stratified fine sand to loamy fine sand	SC-SM, SM, SP-SM	A-2	0	0	100	100	80-95	12-30	0-20	NP-5
Pb: Pits-----	0-60	Gravelly sand	SP-SM, GP-GM, SM, SP	A-1, A-2, A-3	---	0-5	45-100	40-100	0-80	0-40	0-14	NP
Pc: Platte-----	0-9	Loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	85-100	60-95	22-35	4-15
	9-12	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	75-95	45-75	15-30	NP-15
	12-60	Gravelly coarse sand	SM, SP-SM	A-1, A-2, A-3	0	0	70-100	50-95	25-65	0-15	0-10	NP-5
PdD2: Ponca-----	0-4	Silt loam	CL	A-6	0	0	100	100	100	95-100	30-40	10-20
	4-20	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	20-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15

ENGINEERING INDEX PROPERTIES--Continued  
Douglas and Sarpy Counties,  
Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
PdE2: Ponca-----	In											
	0-4	Silt loam	CL	A-6	0	0	100	100	100	95-100	30-40	10-20
	4-20	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	20-60	Silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
Ida-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	5-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
Ra: Riverwash-----	0-6	Sand	SM, SP-SM	A-2, A-3	0	0	100	90-100	65-80	5-35	15-20	NP
	6-60	Stratified sand to very gravelly coarse sand	GP, GW, SP, SW	A-1	0	0-10	50-100	30-90	20-80	3-15	---	NP
Rk: Rock Outcrop---	0-60	Unweathered bedrock			---	---	---	---	---	---	0-14	---
Rn: Ida Variant----	0-6	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
	6-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	95-100	30-40	5-15
Sd: Inavale-----	0-6	Sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	6-60	Fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
Sp: Sarpy-----	0-6	Fine sand	SM, SP, SP-SM	A-2-4, A-3	0	0	100	100	60-80	2-15	---	NP
	6-60	Fine sand	SM, SP, SP-SM	A-2-4, A-3	0	0	100	100	60-80	2-35	---	NP
Ss: Nodaway Variant	0-7	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	95-100	90-100	25-35	5-15
	7-80	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	95-100	90-100	25-35	5-15
StE2: Steinauer-----	0-6	Clay loam	CL	A-6, A-7	0	0-5	90-100	90-100	85-100	55-90	30-50	15-25
	6-18	Clay loam	CH, CL	A-6, A-7	0	0-5	90-100	90-100	90-100	70-90	30-55	12-30
	18-60	Clay loam	CH, CL	A-6, A-7	0	0-5	90-100	90-100	90-100	60-75	25-55	10-30
Wb: Wabash-----	0-15	Silt loam	CL	A-6	0	0	100	100	100	95-100	30-40	10-20
	15-60	Silty clay	CH	A-7	0	0	100	100	100	95-100	52-78	30-55
Wc: Wabash-----	0-7	Silty clay	CH	A-7	0	0	100	100	100	95-100	50-75	30-50
	7-60	Silty clay	CH	A-7	0	0	100	100	100	95-100	52-78	30-55
Wm: Wann-----	0-23	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	95-100	95-100	70-100	30-50	15-25	NP-5
	23-50	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	95-100	75-100	60-100	20-50	15-25	NP-5
	50-60	Stratified sandy clay loam to fine sand	SM	A-2, A-4	0	0	95-100	95-100	70-100	15-40	15-20	NP-3
Wt: Barney-----	0-7	Loam	CL, CL-ML, ML	A-4, A-6	0	0	95-100	90-100	85-95	60-75	20-35	3-15
	7-15	Stratified loam to sand	ML, SM	A-2, A-4	0	0	95-100	90-100	55-80	20-60	10-20	NP-5
	15-60	Gravelly coarse sand	SC-SM, SM, SP, SP-SM	A-1, A-2, A-3	0	0	70-100	50-95	25-65	0-15	0-10	NP-5
zwa: Water-----	---	---	---	---	---	---	---	---	---	---	---	---
zwb: Water-----	---	---	---	---	---	---	---	---	---	---	---	---





## Physical Properties

This table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (ovendry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Saturated hydraulic conductivity refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity (K-sat). The estimates in the table indicate the rate of water movement, in micrometers per second (um/sec), when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In this table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in this table as the K factor (Kw and Kf) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor Kw indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor Kf indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.

PHYSICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

PAGE 2 OF 6

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	in/in	Pct	Pct					

2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.

3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.

4L. Calcareous loams, silt loams, clay loams, and silty clay loams.

4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.

5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.

6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.

7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.

8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

PHYSICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

PAGE 3 OF 6

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Ab: Albaton-----	0-18			22-27	1.30-1.35	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.37	.37	5	4L	86
	18-60			50-60	1.35-1.45	0.00-0.06	0.08-0.12	9.0-25.0	0.0-1.0	.28	.28			
Ac: Albaton-----	0-6			40-60	1.35-1.40	0.06-0.20	0.10-0.14	9.0-25.0	2.0-3.0	.28	.28	5	4	86
	6-60			50-60	1.35-1.45	0.00-0.06	0.08-0.12	9.0-25.0	0.0-1.0	.28	.28			
Af: Alda-----	0-13			5-12	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	2.0-4.0	.20	.20	4	3	86
	13-20			3-10	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.20	.20			
	20-60			0-2	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.15			
Ag: Alda-----	0-13			10-18	1.40-1.60	2.00-6.00	0.20-0.22	0.0-2.9	2.0-4.0	.28	.28	4	3	86
	13-20			3-10	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.20	.20			
	20-60			0-2	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.15			
Ca: Carr-----	0-15			5-15	1.50-1.75	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	4	3	86
	15-60			5-15	1.50-1.75	0.60-6.00	0.13-0.18	0.0-2.9	0.0-0.5	.24	.24			
Cc: Cass-----	0-16			7-17	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-39			5-15	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.28	.28			
	39-60			2-10	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.17	.17			
Cd: Cass-----	0-20			8-18	1.40-1.60	0.57-5.95	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	20-30			8-18	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.20	.20			
	30-48			27-37	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	0.0-0.5	.32	.32			
	48-60			2-8	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.17	.17			
Ce: Cass-----	0-16			8-15	1.40-1.60	2.00-6.00	0.17-0.19	0.0-2.9	1.0-2.0	.32	.32	5	3	86
	16-39			5-15	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.28	.28			
	39-60			2-10	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.17	.17			
Cg: Colo-----	0-23			27-32	1.28-1.32	0.20-0.60	0.21-0.23	3.0-5.9	5.0-7.0	.28	.28	5	7	38
	23-40			30-35	1.25-1.35	0.20-0.60	0.18-0.20	3.0-5.9	3.0-4.0	.28	.28			
	40-60			25-35	1.35-1.45	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.28	.28			
Ck: Colo-----	0-23			27-32	1.28-1.32	0.20-0.60	0.21-0.23	3.0-5.9	5.0-7.0	.28	.28	5	7	38
	23-40			30-35	1.25-1.35	0.20-0.60	0.18-0.20	3.0-5.9	3.0-4.0	.28	.28			
	40-60			25-35	1.35-1.45	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.28	.28			
Kennebec----	0-17			22-27	1.25-1.35	0.60-2.00	0.22-0.24	3.0-5.9	5.0-6.0	.28	.28	5	6	48
	17-60			24-28	1.35-1.40	0.60-2.00	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
Cm: Urban Land---	0-60			---	---	0.06-6.00	0.00-0.00	---	---	---	---	-	8	0
DcE: Dickinson----	0-10			10-18	1.50-1.55	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	10-16			10-18	1.50-1.55	2.00-6.00	0.15-0.17	0.0-2.9	1.0-1.0	.20	.20			
	16-30			10-15	1.45-1.55	2.00-6.00	0.12-0.15	0.0-2.9	0.5-1.0	.24	.24			
	30-60			4-10	1.55-1.65	5.95-19.98	0.08-0.10	0.0-2.9	0.0-0.5	.20	.20			
Ed: Eudora-----	0-13			12-18	1.30-1.50	0.60-2.00	0.20-0.24	0.0-2.9	1.0-4.0	.32	.32	5	5	56
	13-60			5-18	1.35-1.50	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.43	.43			
Ga: Gibbon-----	0-17			2-12	1.60-1.90	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	4	2	134
	17-60			20-27	1.30-1.40	0.60-2.00	0.17-0.19	0.0-2.9	0.5-1.0	.37	.37			
Gb: Gibbon-----	0-16			20-25	1.40-1.60	0.60-2.00	0.21-0.23	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	16-40			20-27	1.30-1.50	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.32	.32			
	40-60			15-25	1.50-1.70	0.57-5.95	0.16-0.20	0.0-2.9	0.5-1.0	.32	.32			
Gc: Gibbon-----	0-16			27-30	1.25-1.35	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.32	.32	5	4L	86
	16-40			27-32	1.50-1.60	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.32	.32			
	40-60			15-30	1.50-1.60	0.20-1.98	0.16-0.20	0.0-2.9	0.5-1.0	.32	.32			
Gs: Gibbon-----	0-20			20-25	1.40-1.60	0.60-2.00	0.21-0.23	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	20-40			27-32	1.50-1.60	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.32	.32			
	40-60			15-30	1.50-1.60	0.20-1.98	0.16-0.20	0.0-2.9	0.5-1.0	.32	.32			
Saltine-----	0-6			15-27	1.30-1.40	0.60-2.00	0.20-0.24	0.0-2.9	0.5-2.0	.37	.37	5	4L	86
	6-15			18-40	1.20-1.30	0.20-0.60	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	15-20			20-45	1.30-1.40	0.20-0.60	0.18-0.20	6.0-8.9	0.0-0.5	.43	.43			
	20-60			20-40	1.40-1.50	0.20-0.60	0.18-0.22	3.0-5.9	0.0-0.5	.43	.43			
Gu: Ida Variant--	0-6			18-27	1.20-1.45	0.60-2.00	0.20-0.22	0.0-2.9	0.5-1.0	.43	.43	4	4L	86
	6-60			18-27	1.40-1.65	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	---			
Ha: Haynie-----	0-9			15-25	1.20-1.35	0.60-2.00	0.18-0.23	0.0-2.9	1.0-3.0	.32	.32	5	4L	86
	9-60			15-18	1.20-1.35	0.60-2.00	0.18-0.23	0.0-2.9	0.0-1.0	.43	.43			
IdD2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
IdE: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	2.0-3.0	.32	.32	5	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
IdE2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

PAGE 4 OF 6

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Im: Inavale-----	0-11			2-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-28			3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.17	.17			
	28-60			3-10	1.50-1.60	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
In: Inavale-----	0-8			2-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	8-25			3-10	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.0-0.5	.17	.17			
	25-60			3-10	1.50-1.60	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
JuB: Judson-----	0-26			24-27	1.30-1.35	0.60-2.00	0.21-0.23	3.0-5.9	4.0-5.0	.28	.28	5	6	48
	26-37			30-35	1.35-1.45	0.40-0.60	0.18-0.20	3.0-5.9	2.0-3.0	.43	.43			
	37-60			25-32	1.35-1.45	0.40-0.60	0.18-0.20	3.0-5.9	0.0-1.0	.43	.43			
Ke: Kennebec-----	0-26			22-27	1.25-1.35	0.60-2.00	0.22-0.24	0.0-2.9	5.0-6.0	.28	.28	5	6	48
	26-60			24-28	1.35-1.40	0.60-2.00	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
Le: Lockton-----	0-21			15-27	1.20-1.40	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	21-28			15-27	1.40-1.60	2.00-6.00	0.12-0.14	3.0-5.9	1.0-3.0	.28	.28			
	28-60			0-5	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.15			
LS: Luton-----	0-11			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	11-27			50-60	1.30-1.35	0.00-0.06	0.01-0.12	9.0-25.0	0.0-1.0	.28	.28			
	27-60			40-60	1.35-1.45	0.00-0.06	0.08-0.12	9.0-25.0	0.0-1.0	.28	.28			
Lt: Luton-----	0-13			35-40	1.30-1.35	0.06-0.20	0.17-0.20	6.0-8.9	3.0-5.0	.32	.32	5	4	86
	13-27			50-60	1.30-1.35	0.00-0.06	0.10-0.12	9.0-25.0	0.0-1.0	.28	.28			
	27-60			40-60	1.35-1.45	0.00-0.06	0.08-0.12	9.0-25.0	0.0-1.0	.28	.28			
Lu: Luton-----	0-17			40-60	1.30-1.35	0.00-0.06	0.12-0.14	9.0-25.0	3.0-5.0	.28	.28	5	4	86
	17-27			50-60	1.30-1.35	0.00-0.06	0.10-0.12	9.0-25.0	0.0-1.0	.28	.28			
	27-60			40-60	1.35-1.45	0.00-0.06	0.08-0.12	9.0-25.0	0.0-1.0	.28	.28			
MaA: Marshall-----	0-16			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.28	.28	5	7	38
	16-39			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.43	.43			
	39-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
MaB: Marshall-----	0-14			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.28	.28	5	7	38
	14-38			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.43	.43			
	38-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
MaC: Marshall-----	0-12			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.28	.28	5	7	38
	12-37			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.43	.43			
	37-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
MaC2: Marshall-----	0-8			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-3.0	.32	.32	5	7	38
	8-32			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	1.0-1.0	.32	.32			
	32-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
MaD: Marshall-----	0-8			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.28	.28	5	7	38
	8-32			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	0.5-1.0	.43	.43			
	32-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
MeD2: Marshall-----	0-8			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-3.0	.32	.32	5	7	38
	8-36			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	1.0-1.0	.32	.32			
	36-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
Ponca-----	0-5			27-32	1.15-1.35	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.32	.32	5	7	38
	5-20			22-32	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	20-60			16-30	1.20-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.43	.43			
MfE: Marshall-----	0-12			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	3.0-4.0	.28	.28	5	7	38
	12-35			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	0.0-1.0	.43	.43			
	35-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
Ponca-----	0-7			27-32	1.15-1.35	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.32	.32	5	7	38
	7-22			22-32	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	22-60			16-30	1.20-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.43	.43			
MfE2: Marshall-----	0-7			27-35	1.25-1.30	0.20-0.60	0.21-0.23	3.0-5.9	2.0-3.0	.32	.32	5	7	38
	7-31			27-34	1.30-1.35	0.20-0.60	0.18-0.20	3.0-5.9	1.0-1.0	.32	.32			
	31-60			22-30	1.30-1.40	0.60-2.00	0.18-0.20	3.0-5.9	0.0-0.5	.43	.43			
Ponca-----	0-4			27-32	1.15-1.35	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.32	.32	5	7	38
	4-20			22-32	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	20-60			16-30	1.20-1.40	0.60-2.00	0.18-0.20	0.0-3.0	0.5-1.0	.43	.43			
MoA: Monona-----	0-12			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	12-35			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	35-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MoB: Monona-----	0-10			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	10-33			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	33-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MoC: Monona-----	0-10			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	10-33			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	33-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

PAGE 5 OF 6

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
MoC2: Pohocco-----	0-7			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	0.5-2.0	.32	.37	5	6	48
	7-30			20-35	1.35-1.45	0.20-0.60	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	30-60			20-27	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MoD: Monona-----	0-10			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	10-33			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	33-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MoD2: Pohocco-----	0-6			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	0.5-2.0	.32	.37	5	6	48
	6-29			20-35	1.35-1.45	0.20-0.60	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	29-60			20-27	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MoE: Monona-----	0-10			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	10-33			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	33-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MsE2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Pohocco-----	0-6			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	0.5-2.0	.32	.37	5	6	48
	6-29			20-35	1.35-1.45	0.20-0.60	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	29-30			20-27	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MsF: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	2.0-3.0	.32	.32	5	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Monona-----	0-8			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	8-31			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	31-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MsF2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Monona-----	0-5			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	5-27			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	27-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
MsG: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	2.0-3.0	.32	.32	5	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Monona-----	0-7			20-27	1.25-1.30	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	7-30			24-28	1.30-1.35	0.60-2.00	0.18-0.20	3.0-5.9	0.5-2.0	.43	.43			
	30-60			18-24	1.35-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
On: Onawa-----	0-7			40-55	1.30-1.35	0.20-0.60	0.12-0.14	6.0-8.9	2.0-3.0	.32	.32	5	4	86
	7-18			50-60	1.30-1.40	0.01-0.06	0.10-0.12	9.0-15.0	0.0-1.0	.32	.32			
	18-60			12-18	1.40-1.50	0.60-5.95	0.16-0.22	0.0-2.9	0.0-0.5	.43	.43			
Pa: Percival-----	0-7			40-60	1.30-1.35	0.01-0.06	0.10-0.12	9.0-25.0	1.0-3.0	.28	.28	4	4	86
	7-16			40-60	1.30-1.35	0.01-0.06	0.10-0.12	9.0-25.0	0.0-2.0	.28	.28			
	16-60			2-12	1.30-1.50	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Pb: Pits-----	0-60			0-8	1.70-2.00	5.95-19.98	0.02-0.09	0.0-2.9	0.0-0.5	.10	.17	2	8	0
Pc: Platte-----	0-9			10-20	1.23-1.45	0.60-2.00	0.20-0.24	0.0-2.9	1.0-3.0	.28	.28	5	4L	86
	9-12			7-18	1.45-1.70	0.57-5.95	0.17-0.19	0.0-2.9	0.0-0.5	.28	.28			
	12-60			0-3	1.65-1.85	1.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
PdD2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Ponca-----	0-4			16-27	1.10-1.30	0.60-2.00	0.21-0.23	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	4-20			22-32	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	20-60			16-30	1.20-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.43	.43			
PdE2: Ida-----	0-5			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	4	4L	86
	5-60			18-25	1.20-1.30	0.60-2.00	0.20-0.22	0.0-2.9	0.0-0.5	.43	.43			
Ponca-----	0-4			16-27	1.10-1.30	0.60-2.00	0.21-0.23	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	4-20			22-32	1.20-1.40	0.20-0.60	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43			
	20-60			16-30	1.20-1.40	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.43	.43			
Ra: Riverwash----	0-6			1-8	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.0-0.1	.15	.15	2	1	220
	6-60			0-5	1.65-1.85	5.95-99.90	0.03-0.05	0.0-2.9	0.0-0.5	.10	.15			
Rk: Rock Outcrop-	0-60			0-0	---	---	0.00-0.00	---	---	---	---	1	8	0
Rn: Ida Variant--	0-6			18-27	1.20-1.45	0.60-2.00	0.20-0.22	0.0-2.9	0.5-1.0	.43	.43	5	4L	86
	6-60			18-27	1.40-1.65	0.60-2.00	0.18-0.20	0.0-2.9	0.0-0.5	.43	---			
Sd: Inavale-----	0-6			1-5	1.50-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	6-60			3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Sp: Sarpy-----	0-6			2-5	1.20-1.50	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	6-60			2-5	1.20-1.50	5.95-19.98	0.05-0.09	0.0-2.9	0.5-1.0	.15	.15			
Ss: Nodaway	0-7			18-27	1.25-1.35	0.60-2.00	0.20-0.23	0.0-2.9	2.0-4.0	.32	.32	5	6	48
Variant-----	7-80			18-30	1.25-1.35	0.60-2.00	0.18-0.20	0.0-2.9	0.5-1.0	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska: Out-of-date

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
StE2: Steinauer----	In	Pct	Pct	Pct	g/cc	in/hr	in/in	Pct	Pct					
	0-6			27-32	1.20-1.35	0.20-0.60	0.19-0.22	3.0-5.9	0.5-2.0	.32	.32	5	4L	86
	6-18			27-32	1.30-1.50	0.20-0.60	0.17-0.19	3.0-5.9	0.5-1.0	.37	.37			
	18-60			24-35	1.30-1.65	0.20-0.60	0.16-0.19	3.0-5.9	0.0-0.5	.37	.37			
Wb: Wabash-----	0-15			20-27	1.35-1.50	0.60-2.00	0.21-0.24	0.0-2.9	2.0-4.0	.28	.28	5	6	48
	15-60			40-60	1.20-1.45	0.00-0.06	0.08-0.12	9.0-25.0	1.0-2.0	.28	.28			
Wc: Wabash-----	0-7			40-46	1.25-1.45	0.00-0.06	0.12-0.14	6.0-8.9	2.0-4.0	.28	.28	5	4	86
	7-60			40-60	1.20-1.45	0.00-0.06	0.08-0.12	9.0-25.0	1.0-2.0	.28	.28			
Wm: Wann-----	0-23			5-15	1.30-1.50	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	23-50			3-18	1.50-1.70	2.00-6.00	0.15-0.17	0.0-2.9	0.5-1.0	.28	.28			
	50-60			3-22	1.35-1.80	2.00-19.98	0.05-0.17	0.0-2.9	0.0-0.5	.15	.15			
Wt: Barney-----	0-7			10-25	1.40-1.50	0.60-2.00	0.20-0.22	0.0-2.9	2.0-4.0	.28	.28	5	8	0
	7-15			3-10	1.60-1.80	1.98-19.98	0.09-0.14	0.0-2.9	0.0-0.5	.17	.17			
	15-60			0-3	1.65-1.85	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
zwa: Water-----	---			---	---	---	---	---	---	---	---	-	---	---
zwb: Water-----	---			---	---	---	---	---	---	---	---	-	---	---





CHEMICAL PROPERTIES OF THE SOILS  
Douglas and Sarpy Counties, Nebraska

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 2 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Ab: Albaton-----	0-18 18-60	20-25 41-50	7.4-8.4 7.4-8.4	5-30 5-30	0 0	0 0	0 0
Ac: Albaton-----	0-6 6-60	36-41 41-50	7.4-8.4 7.4-8.4	5-30 5-30	0 0	0 0	0 0
Af: Alda-----	0-13 13-20 20-60	5.0-20 5.0-10 0.0-5.0	6.6-8.4 7.4-8.4 6.6-8.4	0-10 1-15 0-5	0 0 0	0.0-4.0 0.0-4.0 0.0-4.0	0-9 0-9 0-9
Ag: Alda-----	0-13 13-20 20-60	10-20 5.0-10 0.0-5.0	6.6-8.4 7.4-8.4 6.6-8.4	0-10 1-15 0-5	0 0 0	0.0-4.0 0.0-4.0 0.0-4.0	0-9 0-9 0-9
Ca: Carr-----	0-15 15-60	3.0-11 3.0-10	7.4-8.4 7.4-8.4	1-5 1-10	0 0	0 0	0 0
Cc: Cass-----	0-16 16-39 39-60	6.0-14 4.0-12 1.0-8.0	5.6-7.3 5.6-8.4 6.1-8.4	0 0 0	0 0 0	0 0 0	0 0 0
Cd: Cass-----	0-20 20-30 30-48 48-60	6.0-14 7.0-13 20-25 1.0-8.0	5.6-7.3 6.1-7.8 6.1-7.8 6.1-7.8	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
Ce: Cass-----	0-16 16-39 39-60	7.0-13 4.0-12 1.0-8.0	5.6-7.3 5.6-8.4 6.1-8.4	0 0 0	0 0 0	0 0 0	0 0 0
Cg: Colo-----	0-23 23-40 40-60	36-41 36-41 30-36	5.6-7.3 5.6-7.3 6.1-7.3	0 0 0	0 0 0	0 0 0	--- --- ---
Ck: Colo-----	0-23 23-40 40-60	36-41 36-41 30-36	5.6-7.3 5.6-7.3 6.1-7.3	0 0 0	0 0 0	0 0 0	--- --- ---
Kennebec-----	0-17 17-60	30-36 30-36	5.6-7.3 6.1-7.3	0 0	0 0	0.0-2.0 0.0-2.0	0 0
Cm: Urban Land-----	0-60	---	---	0	0	0	0
DcE: Dickinson-----	0-10 10-16 16-30 30-60	15-20 15-20 15-20 5.0-10	5.6-7.3 5.6-7.3 5.1-6.5 5.1-6.5	--- --- --- ---	--- --- --- ---	--- --- --- ---	--- --- --- ---
Ed: Eudora-----	0-13 13-60	5.0-20 5.0-15	6.1-7.8 6.6-8.4	0 0-10	0 0	0 0	0 0
Ga: Gibbon-----	0-17 17-60	--- 14-20	5.1-5.5 7.4-8.4	0-5 0-5	0 0	0 0	0 0
Gb: Gibbon-----	0-16 16-40 40-60	16-22 14-20 10-18	7.4-8.4 7.4-8.4 7.9-8.4	0-5 5-15 5-15	0 0 0	0.0-2.0 0.0-2.0 0.0-2.0	0 0-5 0-5
Gc: Gibbon-----	0-16 16-40 40-60	20-25 18-26 10-21	7.4-8.4 7.4-8.4 7.9-8.4	0-5 5-15 5-15	0 0 0	0.0-2.0 0.0-2.0 0.0-2.0	0 0-5 0-5
Gs: Gibbon-----	0-20 20-40 40-60	16-22 18-26 10-21	7.4-8.4 7.4-8.4 7.9-8.4	0-5 5-15 5-15	0 0 0	0.0-2.0 0.0-2.0 0.0-2.0	0 0-5 0-5
Saltine-----	0-6 6-15 15-20 20-60	10-20 14-30 14-30 14-30	7.4-9.6 8.5-9.6 7.4-9.6 7.4-9.6	1-10 1-5 1-5 1-5	0 0 0 0	0.0-8.0 4.0-16.0 0.0-4.0 0.0-4.0	0-50 6-99 0-99 0-99

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 3 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Gu:							
Ida Variant-----	0-6	10-20	6.6-8.4	0-10	0	0	0
	6-60	10-20	7.4-8.4	5-15	0	0	0
Ha:							
Haynie-----	0-9	15-20	6.6-8.4	0-25	0	0	0
	9-60	15-20	7.4-8.4	5-30	0	0	0
IdD2:							
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
IdE:							
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
IdE2:							
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
Im:							
Inavale-----	0-11	2.0-8.0	5.6-7.8	0	0	0	0
	11-28	2.0-7.0	5.6-7.8	0	0	0	0
	28-60	2.0-7.0	6.6-8.4	0-5	0	0	0
In:							
Inavale-----	0-8	2.0-8.0	5.6-7.8	0	0	0	0
	8-25	2.0-7.0	5.6-7.8	0	0	0	0
	25-60	2.0-7.0	6.6-8.4	0-5	0	0	0
JuB:							
Judson-----	0-26	25-30	5.6-7.3	0	0	0	0
	26-37	25-30	5.6-7.3	0	0	0	0
	37-60	25-30	6.1-7.8	0-15	0	0	0
Ke:							
Kennebec-----	0-26	30-36	5.6-7.3	0	0	0.0-2.0	0
	26-60	30-36	6.1-7.3	0	0	0.0-2.0	0
Le:							
Lockton-----	0-21	20-30	6.6-7.8	0	0	0	0
	21-28	10-25	6.6-7.8	0	0	0	0
	28-60	0.0-5.0	6.1-7.3	0	0	0	0
Ls:							
Luton-----	0-11	20-25	6.6-7.8	0-15	0	0	0
	11-27	41-200	7.4-8.4	0-15	0	0	0
	27-60	41-200	7.4-8.4	0-25	0	0	0
Lt:							
Luton-----	0-13	41-200	6.6-7.8	0-15	0	0	0
	13-27	41-200	7.4-8.4	0-15	0	0	0
	27-60	41-200	7.4-8.4	0-25	0	0	0
Lu:							
Luton-----	0-17	41-200	6.6-7.8	0-15	0	0	0
	17-27	41-200	7.4-8.4	0-15	0	0	0
	27-60	41-200	7.4-8.4	0-25	0	0	0
MaA:							
Marshall-----	0-16	25-30	5.6-7.3	0	0	0	0
	16-39	25-30	5.6-7.3	0	0	0	0
	39-60	20-25	6.6-7.3	0	0	0	0
MaB:							
Marshall-----	0-14	25-30	5.6-7.3	0	0	0	0
	14-38	25-30	5.6-7.3	0	0	0	0
	38-60	20-25	6.6-7.3	0	0	0	0
MaC:							
Marshall-----	0-12	25-30	5.6-7.3	0	0	0	0
	12-37	25-30	5.6-7.3	0	0	0	0
	37-60	20-25	6.6-7.3	0	0	0	0
MaC2:							
Marshall-----	0-8	25-30	5.6-7.3	0	0	0	0
	8-32	25-30	5.6-7.3	0	0	0	0
	32-60	20-25	6.6-7.3	0	0	0	0
MaD:							
Marshall-----	0-8	25-30	5.6-7.3	0	0	0	0
	8-32	25-30	5.6-7.3	0	0	0	0
	32-60	20-25	6.6-7.3	0	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 4 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
MeD2:							
Marshall-----	0-8	25-30	5.6-7.3	0	0	0	0
	8-36	25-30	5.6-7.3	0	0	0	0
	36-60	20-25	6.6-7.3	0	0	0	0
Ponca-----	0-5	20-25	6.1-7.3	0	0	0	0
	5-20	15-25	6.6-8.4	0-5	0	0	0
	20-60	10-25	7.4-8.4	1-5	0	0	0
MfE:							
Marshall-----	0-12	25-30	5.6-7.3	0	0	0	0
	12-35	25-30	5.6-7.3	0	0	0	0
	35-60	20-25	6.6-7.3	0	0	0	0
Ponca-----	0-7	20-25	6.1-7.3	0	0	0	0
	7-22	15-25	6.6-8.4	0-5	0	0	0
	22-60	10-25	7.4-8.4	1-5	0	0	0
MfE2:							
Marshall-----	0-7	25-30	5.6-7.3	0	0	0	0
	7-31	25-30	5.6-7.3	0	0	0	0
	31-60	20-25	6.6-7.3	0	0	0	0
Ponca-----	0-4	20-25	6.1-7.3	0	0	0	0
	4-20	15-25	6.6-8.4	0-5	0	0	0
	20-60	10-25	7.4-8.4	1-5	0	0	0
MoA:							
Monona-----	0-12	25-30	5.6-7.3	0	0	0	0
	12-35	25-30	6.1-7.3	0	0	0	0
	35-60	20-25	6.6-8.4	0-25	0	0	0
MoB:							
Monona-----	0-10	25-30	5.6-7.3	0	0	0	0
	10-33	25-30	6.1-7.3	0	0	0	0
	33-60	20-25	6.6-8.4	0-25	0	0	0
MoC:							
Monona-----	0-10	25-30	5.6-7.3	0	0	0	0
	10-33	25-30	6.1-7.3	0	0	0	0
	33-60	20-25	6.6-8.4	0-25	0	0	0
MoC2:							
Pohocco-----	0-7	15-25	6.6-7.8	0	0	0	0
	7-30	15-30	6.6-7.8	0	0	0	0
	30-60	15-20	6.6-7.8	0-5	0	0	0
MoD:							
Monona-----	0-10	25-30	5.6-7.3	0	0	0	0
	10-33	25-30	6.1-7.3	0	0	0	0
	33-60	20-25	6.6-8.4	0-25	0	0	0
MoD2:							
Pohocco-----	0-6	15-25	6.6-7.8	0	0	0	0
	6-29	15-30	6.6-7.8	0	0	0	0
	29-60	15-20	6.6-7.8	0-5	0	0	0
MoE:							
Monona-----	0-10	25-30	5.6-7.3	0	0	0	0
	10-33	25-30	6.1-7.3	0	0	0	0
	33-60	20-25	6.6-8.4	0-25	0	0	0
MsE2:							
Pohocco-----	0-6	15-25	6.6-7.8	0	0	0	0
	6-29	15-30	6.6-7.8	0	0	0	0
	29-30	15-20	6.6-7.8	0-5	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
MsF:							
Monona-----	0-8	25-30	5.6-7.3	0	0	0	0
	8-31	25-30	6.1-7.3	0	0	0	0
	31-60	20-25	6.6-8.4	0-25	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
MsF2:							
Monona-----	0-5	25-30	5.6-7.3	0	0	0	0
	5-27	25-30	6.1-7.3	0	0	0	0
	27-60	20-25	6.6-8.4	0-25	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Douglas and Sarpy Counties, Nebraska

PAGE 5 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
MsG:							
Monona-----	0-7	25-30	5.6-7.3	0	0	0	0
	7-30	25-30	6.1-7.3	0	0	0	0
	30-60	20-25	6.6-8.4	0-25	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
On:							
Onawa-----	0-7	36-41	7.4-8.4	5-30	0	0	0
	7-18	41-50	7.4-8.4	5-30	0	0	0
	18-60	15-20	7.4-8.4	5-30	0	0	0
Pa:							
Percival-----	0-7	36-41	7.4-8.4	0-15	0	0	0
	7-16	35-40	7.4-8.4	0-25	0	0	0
	16-60	5.0-15	7.4-8.4	0-25	0	0	0
Pb:							
Pits-----	0-60	0.0-5.0	6.6-8.4	0	0	0	0
Pc:							
Platte-----	0-9	5.0-20	6.6-8.4	0-10	0	0.0-2.0	0
	9-12	5.0-20	6.6-8.4	0-10	0	0.0-2.0	0
	12-60	0.0-5.0	6.6-8.4	0-5	0	0.0-2.0	0
PdD2:							
Ponca-----	0-4	15-25	6.1-7.3	0	0	0	0
	4-20	15-25	6.6-8.4	0-5	0	0	0
	20-60	10-25	7.4-8.4	1-5	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
PdE2:							
Ponca-----	0-4	15-25	6.1-7.3	0	0	0	0
	4-20	15-25	6.6-8.4	0-5	0	0	0
	20-60	10-25	7.4-8.4	1-5	0	0	0
Ida-----	0-5	20-25	6.6-8.4	0-25	0	0	0
	5-60	20-25	7.4-8.4	5-30	0	0	0
Ra:							
Riverwash-----	0-6	1.0-10	7.4-8.4	0-5	0	0	0
	6-60	0.0-5.0	7.4-8.4	0-5	0	0	0
Rk:							
Rock Outcrop----	0-60	---	---	---	---	0	---
Rn:							
Ida Variant-----	0-6	10-20	6.6-8.4	0-10	0	0	0
	6-60	10-20	7.4-8.4	5-15	0	0	0
Sd:							
Inavale-----	0-6	1.0-4.0	5.6-7.8	0	0	0	0
	6-60	2.0-7.0	6.6-8.4	0-5	0	0	0
Sp:							
Sarpy-----	0-6	2.0-6.0	6.6-8.4	1-2	---	---	---
	6-60	2.0-6.0	6.6-8.4	1-2	---	---	---
Ss:							
Nodaway Variant--	0-7	15-30	6.1-7.3	0	0	0.0-2.0	0
	7-80	10-25	6.1-7.3	0	0	0.0-2.0	0
StE2:							
Steinauer-----	0-6	15-25	7.4-8.4	1-10	0	0	0
	6-18	15-25	7.4-8.4	5-15	0	0	0
	18-60	15-25	7.9-8.4	5-20	0	0	0
Wb:							
Wabash-----	0-15	15-20	5.1-7.3	0	0	0	0
	15-60	28-42	5.1-7.8	0	0	0	0
Wc:							
Wabash-----	0-7	30-35	5.1-7.3	0	0	0	0
	7-60	28-42	5.1-7.8	0	0	0	0
Wm:							
Wann-----	0-23	4.0-14	6.6-8.4	0-5	0	0.0-2.0	0-5
	23-50	2.0-14	7.4-9.0	0-5	0	0.0-2.0	0-10
	50-60	2.0-16	7.4-9.0	0-5	0	0.0-2.0	0-10
Wt:							
Barney-----	0-7	9.0-20	6.6-8.4	0-5	0	0	0
	7-15	2.0-7.0	6.6-8.4	0-5	0	0	0
	15-60	0.0-2.0	6.6-7.8	0	0	0	0
zwa:							
Water-----	---	---	---	---	---	---	---
zwb:							
Water-----	---	---	---	---	---	---	---



WATER FEATURES  
Douglas and Sarpy Counties, Nebraska

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Ab: Albaton-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		March	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		April	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		May	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		June	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		July	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		December	0.0-1.5	>6.0	---	---	---	---	None
Ac: Albaton-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		March	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		April	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		May	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		June	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		July	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	0.0-1.5	>6.0	---	---	---	Brief	Occasional
		December	0.0-1.5	>6.0	---	---	---	---	None
Af: Alda-----	C	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Ag: Alda-----	C	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Ca: Carr-----	B	March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
Cc: Cass-----	B	March	---	---	---	---	---	Brief	Rare
		April	---	---	---	---	---	Brief	Rare
		May	---	---	---	---	---	Brief	Rare
		June	---	---	---	---	---	Brief	Rare
		July	---	---	---	---	---	Brief	Rare
Cd: Cass-----	B	January	2.5-3.5	2.5-3.5	---	---	---	---	None
		February	2.5-3.5	2.5-3.5	---	---	---	---	None
		March	2.5-3.5	2.5-3.5	---	---	---	Brief	Rare
		April	2.5-3.5	2.5-3.5	---	---	---	Brief	Rare
		May	2.5-3.5	2.5-3.5	---	---	---	Brief	Rare
		June	2.5-3.5	2.5-3.5	---	---	---	Brief	Rare
		July	---	---	---	---	---	Brief	Rare
		November	2.5-3.5	2.5-3.5	---	---	---	---	None
		December	2.5-3.5	2.5-3.5	---	---	---	---	None
Ce: Cass-----	B	March	---	---	---	---	---	Brief	Rare
		April	---	---	---	---	---	Brief	Rare
		May	---	---	---	---	---	Brief	Rare
		June	---	---	---	---	---	Brief	Rare
		July	---	---	---	---	---	Brief	Rare



(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Cg: Colo-----	B/D		Ft	Ft	Ft				
		January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Long	Occasional
		March	0.0-1.5	>6.0	---	---	---	Long	Occasional
		April	0.0-1.5	>6.0	---	---	---	Long	Occasional
		May	0.0-1.5	>6.0	---	---	---	Long	Occasional
		June	0.0-1.5	>6.0	---	---	---	Long	Occasional
		July	0.0-1.5	>6.0	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	0.0-1.5	>6.0	---	---	---	Long	Occasional
		December	0.0-1.5	>6.0	---	---	---	---	None
Ck: Colo-----	B/D								
		January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Long	Occasional
		March	0.0-1.5	>6.0	---	---	---	Long	Occasional
		April	0.0-1.5	>6.0	---	---	---	Long	Occasional
		May	0.0-1.5	>6.0	---	---	---	Long	Occasional
		June	0.0-1.5	>6.0	---	---	---	Long	Occasional
		July	0.0-1.5	>6.0	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	0.0-1.5	>6.0	---	---	---	Long	Occasional
		December	0.0-1.5	>6.0	---	---	---	---	None
Kennebec-----	B								
		January	3.0-6.0	>6.0	---	---	---	---	None
		February	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		March	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		April	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		May	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		June	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		July	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		December	3.0-6.0	>6.0	---	---	---	---	None
Cm: Urban Land-----	D								
			---	---	---	---	---	---	---
DcE: Dickinson-----	B								
			---	---	---	---	---	---	---
Ed: Eudora-----	B								
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
			---	---	---	---	---	---	---
Ga: Gibbon-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Gb: Gibbon-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Gc:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Gibbon-----	B		Ft	Ft	Ft				
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Gs: Gibbon-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Saltine-----	C								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		July	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		December	1.5-3.0	>6.0	---	---	---	---	None
Gu: Ida Variant-----	B								
			---	---	---	---	---	---	---
Ha: Haynie-----	B								
			---	---	---	---	---	---	---
IdD2: Ida-----	B								
			---	---	---	---	---	---	---
IdE: Ida-----	B								
			---	---	---	---	---	---	---
IdE2: Ida-----	B								
			---	---	---	---	---	---	---
Im: Inavale-----	A								
			---	---	---	---	---	---	---
In: Inavale-----	A								
			---	---	---	---	---	---	---
JuB: Judson-----	B								
			---	---	---	---	---	---	---
Ke: Kennebec-----	B								
		January	3.0-6.0	>6.0	---	---	---	---	None
		February	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		March	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		April	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		May	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		June	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		July	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	3.0-6.0	>6.0	---	---	---	Brief	Occasional
		December	3.0-6.0	>6.0	---	---	---	---	None
Le: Lockton-----	B								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Ls:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Luton-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	Brief	Rare
		April	0.0-1.5	>6.0	---	---	---	Brief	Rare
		May	0.0-1.5	>6.0	---	---	---	Brief	Rare
		June	0.0-1.5	>6.0	---	---	---	Brief	Rare
		July	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Lt: Luton-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	Brief	Rare
		April	0.0-1.5	>6.0	---	---	---	Brief	Rare
		May	0.0-1.5	>6.0	---	---	---	Brief	Rare
		June	0.0-1.5	>6.0	---	---	---	Brief	Rare
		July	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Lu: Luton-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	Brief	Rare
		April	0.0-1.5	>6.0	---	---	---	Brief	Rare
		May	0.0-1.5	>6.0	---	---	---	Brief	Rare
		June	0.0-1.5	>6.0	---	---	---	Brief	Rare
		July	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
MaA: Marshall-----	B	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	Brief	Rare
		April	0.0-1.5	>6.0	---	---	---	Brief	Rare
		May	0.0-1.5	>6.0	---	---	---	Brief	Rare
		June	0.0-1.5	>6.0	---	---	---	Brief	Rare
		July	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
MaB: Marshall-----	B		---	---	---	---	---	---	---
MaC: Marshall-----	B		---	---	---	---	---	---	---
MaC2: Marshall-----	B		---	---	---	---	---	---	---
MaD: Marshall-----	B		---	---	---	---	---	---	---
MeD2: Marshall-----	B		---	---	---	---	---	---	---
Ponca-----	B		---	---	---	---	---	---	---
MfE: Marshall-----	B		---	---	---	---	---	---	---
Ponca-----	B		---	---	---	---	---	---	---
MfE2: Marshall-----	B		---	---	---	---	---	---	---
Ponca-----	B		---	---	---	---	---	---	---
MoA: Monona-----	B		---	---	---	---	---	---	---
MoB: Monona-----	B		---	---	---	---	---	---	---
MoC: Monona-----	B		---	---	---	---	---	---	---
MoC2: Pohocco-----	B		---	---	---	---	---	---	---
MoD: Monona-----	B		---	---	---	---	---	---	---
MoD2: Pohocco-----	B		---	---	---	---	---	---	---
MoE: Monona-----	B		---	---	---	---	---	---	---
MsE2:			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Ida-----	B		Ft	Ft	Ft				
Pohocco-----	B		---	---	---	---	---	---	---
MsF: Ida-----	B		---	---	---	---	---	---	---
Monona-----	B		---	---	---	---	---	---	---
MsF2: Ida-----	B		---	---	---	---	---	---	---
Monona-----	B		---	---	---	---	---	---	---
MsG: Ida-----	B		---	---	---	---	---	---	---
Monona-----	B		---	---	---	---	---	---	---
On: Onawa-----	D		---	---	---	---	---	---	---
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		July	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		December	1.5-3.0	>6.0	---	---	---	---	None
Pa: Percival-----	C		---	---	---	---	---	---	None
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		March	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		June	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		July	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	Very brief	Occasional
		December	1.5-3.0	>6.0	---	---	---	---	None
Pb: Pits-----	A		---	---	---	---	---	---	---
Pc: Platte-----	B		---	---	---	---	---	---	---
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
PdD2: Ida-----	B		---	---	---	---	---	---	---
Ponca-----	B		---	---	---	---	---	---	---
PdE2: Ida-----	B		---	---	---	---	---	---	---
Ponca-----	B		---	---	---	---	---	---	---
Ra: Riverwash-----	A		---	---	---	---	---	---	---
		January	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		February	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		March	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		April	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		May	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		June	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		July	---	---	---	---	---	Very long	Frequent
		October	---	---	---	---	---	Very long	Frequent
		November	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
		December	0.0	>6.0	0.0-2.0	Very long	---	Very long	Frequent
Rk:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Rock Outcrop-----	D		Ft	Ft	Ft				
Rn: Ida Variant-----	B		---	---	---	---	---	---	---
Sd: Inavale-----	A		---	---	---	---	---	---	---
Sp: Sarpy-----	A	January	---	---	---	---	---	Very brief	Frequent
		February	---	---	---	---	---	Very brief	Frequent
		March	---	---	---	---	---	Very brief	Frequent
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	---	---
		September	---	---	---	---	---	---	---
		October	---	---	---	---	---	---	---
		November	---	---	---	---	---	---	---
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
Ss: Nodaway Variant-----	B	February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
StE2: Steinauer-----	B	March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
Wb: Wabash-----	D	April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
Wc: Wabash-----	D	May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
Wm: Wann-----	B	June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
Wt: Barney-----	D	July	---	---	---	---	---	Long	Occasional
		August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
		October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional
		December	---	---	---	---	---	Long	Occasional
		January	---	---	---	---	---	Long	Occasional
		February	---	---	---	---	---	Long	Occasional
		March	---	---	---	---	---	Long	Occasional
		April	---	---	---	---	---	Long	Occasional
		May	---	---	---	---	---	Long	Occasional
		June	---	---	---	---	---	Long	Occasional
		July	---	---	---	---	---	Long	Occasional
zwa: Water-----	---	August	---	---	---	---	---	Long	Occasional
		September	---	---	---	---	---	Long	Occasional
zwb: Water-----	---	October	---	---	---	---	---	Long	Occasional
		November	---	---	---	---	---	Long	Occasional

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft ---	Ft ---	Ft ---	---	---	---	---



SOIL FEATURES  
Douglas and Sarpy Counties, Nebraska

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.



Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
		In	In				
Ab: Albaton-----	---	---	---	---	Moderate	High	Low
Ac: Albaton-----	---	---	---	---	Moderate	High	Low
Af: Alda-----	---	---	---	---	High	Moderate	Low
Ag: Alda-----	---	---	---	---	High	Moderate	Low
Ca: Carr-----	---	---	---	---	Low	Low	Low
Cc: Cass-----	---	---	---	---	Moderate	Moderate	Low
Cd: Cass-----	---	---	---	---	Moderate	Moderate	Low
Ce: Cass-----	---	---	---	---	Moderate	Moderate	Low
Cg: Colo-----	---	---	---	---	High	High	Moderate
Ck: Colo-----	---	---	---	---	High	High	Moderate
Cm: Kennebec-----	---	---	---	---	High	Moderate	Low
Urban Land-----	---	---	---	---	None	---	---
DcE: Dickinson-----	---	---	---	---	Moderate	Low	Moderate
Ed: Eudora-----	---	---	---	---	High	Low	Low
Ga: Gibbon-----	---	---	---	---	High	Moderate	Low
Gb: Gibbon-----	---	---	---	---	High	High	Low
Gc: Gibbon-----	---	---	---	---	High	High	Low
Gs: Gibbon-----	---	---	---	---	High	High	Low
Gu: Saltine-----	---	---	---	---	High	High	High
Ida Variant-----	---	---	---	---	High	Low	Low
Ha: Haynie-----	---	---	---	---	High	Low	Low
IdD2: Ida-----	---	---	---	---	High	Low	Low
IdE: Ida-----	---	---	---	---	High	Low	Low
IdE2: Ida-----	---	---	---	---	High	Low	Low
Im: Inavale-----	---	---	---	---	Low	Moderate	Low
In: Inavale-----	---	---	---	---	Low	Moderate	Low
JuB: Judson-----	---	---	---	---	High	Moderate	Low
Ke: Kennebec-----	---	---	---	---	High	Moderate	Low
Le: Lockton-----	---	---	---	---	High	Moderate	Low
Ls: Luton-----	---	---	---	---	Moderate	High	Low
Lt: Luton-----	---	---	---	---	Moderate	High	Low
Lu: Luton-----	---	---	---	---	Moderate	High	Low
MaA: Marshall-----	---	---	---	---	High	Moderate	Moderate
MaB: Marshall-----	---	---	---	---	High	Moderate	Moderate
MaC: Marshall-----	---	---	---	---	High	Moderate	Moderate
MaC2: Marshall-----	---	---	---	---	High	Moderate	Moderate
MaD: Marshall-----	---	---	---	---	High	Moderate	Moderate
MeD2: Marshall-----	---	---	---	---	High	Moderate	Moderate
MfE: Ponca-----	---	---	---	---	High	Moderate	Low
MfE2: Marshall-----	---	---	---	---	High	Moderate	Moderate
MoA: Monona-----	---	---	---	---	High	Low	Low
MoB: Monona-----	---	---	---	---	High	Low	Low
MoC: Monona-----	---	---	---	---	High	Low	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
		In	In				
MoC2: Pohocco-----	---	---	---	---	High	Moderate	Low
MoD: Monona-----	---	---	---	---	High	Low	Low
MoD2: Pohocco-----	---	---	---	---	High	Moderate	Low
MoE: Monona-----	---	---	---	---	High	Low	Low
MsE2: Ida-----	---	---	---	---	High	Low	Low
Pohocco-----	---	---	---	---	High	Moderate	Low
MsF: Ida-----	---	---	---	---	High	Low	Low
Monona-----	---	---	---	---	High	Low	Low
MsF2: Ida-----	---	---	---	---	High	Low	Low
Monona-----	---	---	---	---	High	Low	Low
MsG: Ida-----	---	---	---	---	High	Low	Low
Monona-----	---	---	---	---	High	Low	Low
On: Onawa-----	---	---	---	---	High	High	Low
Pa: Percival-----	---	---	---	---	Moderate	High	Low
Pb: Pits-----	---	---	---	---	Low	Low	Low
Pc: Platte-----	---	---	---	---	Moderate	High	Moderate
PdD2: Ida-----	---	---	---	---	High	Low	Low
Ponca-----	---	---	---	---	High	Moderate	Low
PdE2: Ida-----	---	---	---	---	High	Low	Low
Ponca-----	---	---	---	---	High	Moderate	Low
Ra: Riverwash-----	---	---	---	---	Low	Moderate	Moderate
Rk: Rock Outcrop----	0-0	Bedrock (paralithic)	---	---	None	---	---
Rn: Ida Variant-----	---	---	---	---	High	Low	Low
Sd: Inavale-----	---	---	---	---	Low	Moderate	Low
Sp: Sarpy-----	---	---	---	---	Low	Low	Low
Ss: Nodaway Variant-----	---	---	---	---	High	Moderate	Low
StE2: Steinauer-----	---	---	---	---	Moderate	High	Low
Wb: Wabash-----	---	---	---	---	Moderate	High	Moderate
Wc: Wabash-----	---	---	---	---	Moderate	High	Moderate
Wm: Wann-----	---	---	---	---	High	Moderate	Low
Wt: Barney-----	---	---	---	---	Moderate	High	Low
zwa: Water-----	---	---	---	---	---	---	---
zwb: Water-----	---	---	---	---	---	---	---



WATER MANAGEMENT  
Douglas and Sarpy Counties, Nebraska

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Ab: Albaton-----	Limitation: flooding percs slowly	Limitation: percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
Ac: Albaton-----	Limitation: flooding percs slowly	Limitation: percs slowly slow intake wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
Af: Alda-----	Limitation: flooding frost action cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Favorable
Ag: Alda-----	Limitation: flooding frost action cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Favorable
Ca: Carr-----	Limitation: deep to water	Limitation: flooding rooting depth	Limitation: soil blowing	Limitation: rooting depth
Cc: Cass-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Cd: Cass-----	Favorable	Limitation: wetness soil blowing	Limitation: wetness soil blowing	Favorable
Ce: Cass-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Cg: Colo-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Limitation: wetness
Ck: Colo-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Limitation: wetness
Kennebec-----	Limitation: deep to water	Limitation: flooding	Favorable	Favorable
Cm: Urban Land-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope
DcE: Dickinson-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope too sandy soil blowing	Limitation: slope
Ed: Eudora-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Ga: Gibbon-----	Limitation: flooding frost action	Limitation: fast intake wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily rooting depth
Gb: Gibbon-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Favorable
Gc: Gibbon-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Favorable
Gs: Gibbon-----	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Favorable
Saltine-----	Limitation: excess salt flooding frost action	Limitation: erodes easily flooding wetness	Limitation: erodes easily wetness	Limitation: erodes easily excess sodium excess salt
Gu: Ida Variant-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ha: Haynie-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
IdD2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
IdE: Ida-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
IdE2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Im: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
In: Inavale-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
JuB: Judson-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
Ke: Kennebec-----	Limitation: deep to water	Limitation: flooding	Favorable	Favorable
Le: Lockton-----	Limitation: deep to water	Favorable	Limitation: too sandy	Favorable
Ls: Luton-----	Limitation: flooding percs slowly	Limitation: percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
Lt: Luton-----	Limitation: flooding percs slowly	Limitation: percs slowly wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
Lu: Luton-----	Limitation: flooding percs slowly	Limitation: percs slowly slow intake wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
MaA: Marshall-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
MaB: Marshall-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
MaC: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
MaC2: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
MaD: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MeD2: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ponca-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MfE: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ponca-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MfE2: Marshall-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ponca-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MoA: Monona-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
MoB: Monona-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
MoC: Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
MoC2: Pohocco-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily	Limitation: erodes easily
MoD: Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MoD2: Pohocco-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MoE: Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MsE2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Pohocco-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MsF: Ida-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MsF2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
MsG: Ida-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Monona-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
On: Onawa-----	Limitation: flooding frost action percs slowly	Limitation: percs slowly slow intake wetness	Limitation: erodes easily wetness	Limitation: erodes easily percs slowly
Pa: Percival-----	Limitation: flooding percs slowly cutbanks cave	Limitation: slow intake wetness droughty	Limitation: too sandy wetness	Limitation: percs slowly droughty
Pb: Pits-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy	Limitation: rooting depth slope droughty
Pc: Platte-----	Limitation: flooding cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
PdD2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ponca-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
PdE2: Ida-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Ponca-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Ra: Riverwash-----	Limitation: flooding ponding cutbanks cave	Limitation: fast intake ponding droughty	Limitation: too sandy ponding	Limitation: rooting depth wetness droughty
Rk: Rock Outcrop----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
Rn: Ida Variant-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Sd: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Sp: Sarpy-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Ss: Nodaway Variant-	Limitation: flooding frost action	Limitation: flooding wetness	Limitation: wetness	Favorable
StE2: Steinauer-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily slope	Limitation: erodes easily slope
Wb: Wabash-----	Limitation: flooding percs slowly	Limitation: wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
Wc: Wabash-----	Limitation: flooding percs slowly	Limitation: slow intake wetness droughty	Limitation: percs slowly wetness	Limitation: percs slowly wetness droughty
Wm: Wann-----	Limitation: flooding frost action	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Favorable
Wt: Barney-----	Limitation: flooding cutbanks cave	Limitation: rooting depth wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
zwa: Water-----	---	---	---	---
zwb: Water-----	---	---	---	---



WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Ac: Albaton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Very limited Slow refill Cutbanks cave	1.00 0.10
Af: Alda-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage Piping	0.95 0.14 0.02	Very limited Cutbanks cave Deep to water	1.00 0.02
Ag: Alda-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage Piping	0.95 0.14 0.02	Very limited Cutbanks cave Deep to water	1.00 0.02
Ca: Carr-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.00	Very limited Deep to water	1.00
Cc: Cass-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Deep to water	1.00
Cd: Cass-----	100	Very limited Seepage	1.00	Very limited Piping Depth to saturated zone Seepage	1.00 0.43 0.22	Very limited Cutbanks cave Deep to water	1.00 0.25
Ce: Cass-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Deep to water	1.00
Cg: Colo-----	100	Somewhat limited Seepage	0.04	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10
Ck: Colo-----	50	Somewhat limited Seepage	0.04	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10
Kennebec-----	50	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.99	Somewhat limited Deep to water Slow refill Cutbanks cave	0.96 0.30 0.10
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Very limited Seepage Slope	1.00 0.02	Somewhat limited Seepage	0.72	Very limited Deep to water	1.00
Ed: Eudora-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Ga: Gibbon-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.97	Somewhat limited Slow refill	0.30

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Gb: Gibbon-----	100	Very limited Seepage	1.00	Depth to saturated zone	0.95	Cutbanks cave	0.10
				Seepage	0.16	Deep to water	0.02
				Very limited Piping	1.00	Somewhat limited Cutbanks cave	0.10
				Depth to saturated zone	0.95	Deep to water	0.02
Gc: Gibbon-----	100	Somewhat limited Seepage	0.12	Somewhat limited Depth to saturated zone	0.95	Somewhat limited Slow refill	0.88
				Piping	0.44	Cutbanks cave	0.10
						Deep to water	0.02
Gs: Gibbon-----	65	Somewhat limited Seepage	0.70	Somewhat limited Depth to saturated zone	0.95	Somewhat limited Slow refill	0.88
				Piping	0.79	Cutbanks cave	0.10
						Deep to water	0.02
Saltine-----	35	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Somewhat limited Slow refill	0.96
				Depth to saturated zone	0.95	Salty water	0.50
				Salinity	0.12	Cutbanks cave	0.10
Gu: Ida Variant-----	100	Very limited Slope Seepage	1.00 0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Ha: Haynie-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
IdD2: Ida-----	100	Somewhat limited Seepage Slope	0.70 0.00	Very limited Piping	1.00	Very limited Deep to water	1.00
IdE: Ida-----	100	Somewhat limited Seepage Slope	0.70 0.24	Very limited Piping	1.00	Very limited Deep to water	1.00
IdE2: Ida-----	100	Somewhat limited Seepage Slope	0.70 0.24	Very limited Piping	1.00	Very limited Deep to water	1.00
Im: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.33	Very limited Deep to water	1.00
In: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.33	Very limited Deep to water	1.00
JuB: Judson-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.60	Very limited Deep to water	1.00
Ke: Kennebec-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.95	Somewhat limited Deep to water	0.96
Le: Lockton-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95	Very limited Cutbanks cave	1.00
					0.28	Deep to water	0.02

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ls: Luton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Lt: Luton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Very limited Slow refill Cutbanks cave	1.00 0.10
Lu: Luton-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Very limited Slow refill Cutbanks cave	1.00 0.10
MaA: Marshall-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
MaB: Marshall-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
MaC: Marshall-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
MaC2: Marshall-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
MaD: Marshall-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
MeD2: Marshall-----	70	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Ponca-----	30	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
MfE: Marshall-----	50	Somewhat limited Seepage Slope	0.70 0.02	Not limited		Very limited Deep to water	1.00
Ponca-----	50	Somewhat limited Seepage Slope	0.70 0.02	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
MfE2: Marshall-----	50	Somewhat limited Seepage Slope	0.70 0.02	Not limited		Very limited Deep to water	1.00
Ponca-----	50	Somewhat limited Seepage Slope	0.70 0.02	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
MoA: Monona-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.28	Very limited Deep to water	1.00
MoB: Monona-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.29	Very limited Deep to water	1.00
MoC: Monona-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.29	Very limited Deep to water	1.00
MoC2: Pohocco-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.82	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MoD: Monona-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.29	Very limited Deep to water	1.00
MoD2: Pohocco-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.83	Very limited Deep to water	1.00
MoE: Monona-----	100	Somewhat limited Seepage Slope	0.70 0.02	Somewhat limited Piping	0.29	Very limited Deep to water	1.00
MsE2: Ida-----	50	Somewhat limited Seepage Slope	0.70 0.04	Very limited Piping	1.00	Very limited Deep to water	1.00
Pohocco-----	50	Somewhat limited Seepage Slope	0.70 0.02	Somewhat limited Piping	0.16	Very limited Deep to water	1.00
MsF: Ida-----	50	Somewhat limited Seepage Slope	0.70 0.28	Very limited Piping	1.00	Very limited Deep to water	1.00
Monona-----	50	Somewhat limited Seepage Slope	0.70 0.24	Somewhat limited Piping	0.31	Very limited Deep to water	1.00
MsF2: Ida-----	50	Somewhat limited Seepage Slope	0.70 0.28	Very limited Piping	1.00	Very limited Deep to water	1.00
Monona-----	50	Somewhat limited Seepage Slope	0.70 0.24	Somewhat limited Piping	0.34	Very limited Deep to water	1.00
MsG: Ida-----	50	Somewhat limited Slope Seepage	0.97 0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Monona-----	50	Somewhat limited Slope Seepage	0.72 0.70	Somewhat limited Piping	0.32	Very limited Deep to water	1.00
On: Onawa-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Piping	0.95 0.03	Somewhat limited Cutbanks cave Deep to water	0.10 0.02
Pa: Percival-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.19	Very limited Cutbanks cave Deep to water	1.00 0.02
Pb: Pits-----	100	Not rated		Not rated		Not rated	
Pc: Platte-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.16	Very limited Cutbanks cave Deep to water	1.00 0.02
PdD2: Ida-----	50	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Ponca-----	50	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
PdE2: Ida-----	50	Somewhat limited Seepage Slope	0.70 0.04	Very limited Piping	1.00	Very limited Deep to water	1.00
Ponca-----	50	Somewhat limited Seepage Slope	0.70 0.02	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
Ra: Riverwash-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.22	Very limited Cutbanks cave	1.00
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	
Rn: Ida Variant-----	100	Somewhat limited Slope Seepage	0.97 0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Sd: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.33	Very limited Deep to water	1.00
Sp: Sarpy-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.96	Very limited Deep to water	1.00
Ss: Nodaway Variant----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Somewhat limited Deep to water Slow refill Cutbanks cave	0.96 0.30 0.10
StE2: Steinauer-----	100	Somewhat limited Slope Seepage	0.15 0.05	Not limited		Very limited Deep to water	1.00
Wb: Wabash-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Wc: Wabash-----	100	Not limited		Very limited Depth to saturated zone Hard to pack	1.00 1.00	Very limited Slow refill Cutbanks cave	1.00 0.10
Wm: Wann-----	100	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage Piping	0.84 0.10 0.02	Very limited Cutbanks cave Deep to water	1.00 0.07
Wt: Barney-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.16	Very limited Cutbanks cave	1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

WATER MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit						



SANITARY FACILITIES  
Douglas and Sarpy Counties, Nebraska

### Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.



SANITARY FACILITIES  
Douglas and Sarpy Counties, Nebraska

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Ac: Albaton-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Af: Alda-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Ag: Alda-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Ca: Carr-----	100	Very limited Flooding	1.00	Very limited Flooding Seepage	1.00 1.00
Cc: Cass-----	100	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40
Cd: Cass-----	100	Very limited Depth to saturated zone Restricted permeability Filtering capacity Flooding	1.00 1.00 1.00 0.40	Very limited Seepage Depth to saturated zone Flooding	1.00 1.00 0.40
Ce: Cass-----	100	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40
Cg: Colo-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Ck: Colo-----	50	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Kennebec-----	50	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 0.84 0.50	Very limited Flooding Seepage Depth to saturated zone	1.00 0.50 0.17
Cm: Urban Land-----	100	Not rated		Not rated	
DcE: Dickinson-----	100	Very limited Filtering capacity Slope	1.00 0.96	Very limited Slope Seepage	1.00 1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ed: Eudora-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage Flooding	0.50 0.40
Ga: Gibbon-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.50
Gb: Gibbon-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00
Gc: Gibbon-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.00
Gs: Gibbon-----	65	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.00
Saltine-----	35	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Gu: Ida Variant-----	100	Very limited Slope Restricted permeability	1.00 0.50	Very limited Slope Seepage	1.00 0.50
Ha: Haynie-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
IdD2: Ida-----	100	Somewhat limited Slope Restricted permeability	0.63 0.50	Very limited Slope Seepage	1.00 0.50
IdE: Ida-----	100	Very limited Slope Restricted permeability	1.00 0.50	Very limited Slope Seepage	1.00 0.50
IdE2: Ida-----	100	Very limited Slope Restricted permeability	1.00 0.50	Very limited Slope Seepage	1.00 0.50
Im: Inavale-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
In: Inavale-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.67
JuB: Judson-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
Ke: Kennebec-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Le: Lockton-----	100	Depth to saturated zone	0.84	Seepage	0.50
		Restricted permeability	0.50	Depth to saturated zone	0.17
		Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
Ls: Luton-----	100	Filtering zone capacity	1.00	Depth to saturated zone	1.00
		Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Flooding	0.40
Lt: Luton-----	100	Flooding	0.40		
		Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Flooding	0.40
Lu: Luton-----	100	Flooding	0.40		
		Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Flooding	0.40
MaA: Marshall-----	100	Flooding	0.40		
MaB: Marshall-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
		Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
MaC: Marshall-----	100			Slope	0.00
		Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
MaC2: Marshall-----	100			Seepage	0.50
		Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
MaD: Marshall-----	100			Seepage	0.50
		Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
MeD2: Marshall-----	70	Slope	0.04	Seepage	0.50
		Very limited Restricted permeability	1.00	Very limited Slope	1.00
Ponca-----	30	Slope	0.04	Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
MfE: Marshall-----	50	Slope	0.04	Seepage	0.50
		Very limited Restricted permeability	1.00	Very limited Slope	1.00
Ponca-----	50	Slope	0.96	Seepage	0.50
		Somewhat limited Slope	0.96	Very limited Slope	1.00
MfE2: Marshall-----	50	Restricted permeability	0.50	Seepage	0.50
		Very limited Restricted permeability	1.00	Very limited Slope	1.00
Ponca-----	50	Slope	0.96	Seepage	0.50
		Somewhat limited		Very limited	

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
MoA: Monona-----	100	Slope	0.96	Slope	1.00
		Restricted permeability	0.50	Seepage	0.50
MoB: Monona-----	100	Somewhat limited	0.50	Somewhat limited	0.50
		Restricted permeability		Seepage	
MoC: Monona-----	100	Somewhat limited	0.50	Somewhat limited	0.50
		Restricted permeability		Seepage	
MoC2: Pohocco-----	100	Somewhat limited	0.50	Slope	0.00
		Restricted permeability		Seepage	
MoD: Monona-----	100	Somewhat limited	0.50	Somewhat limited	0.67
		Restricted permeability		Slope	
MoD2: Pohocco-----	100	Very limited	1.00	Seepage	0.50
		Restricted permeability		Slope	
MoE: Monona-----	100	Somewhat limited	0.50	Very limited	1.00
		Slope		Slope	
MsE2: Ida-----	50	Restricted permeability	0.50	Seepage	0.50
		Slope		Slope	
Pohocco-----	50	Very limited	1.00	Very limited	1.00
		Restricted permeability		Slope	
MsF: Ida-----	50	Slope	0.96	Seepage	0.50
		Restricted permeability		Slope	
Monona-----	50	Very limited	1.00	Very limited	1.00
		Slope		Slope	
MsF2: Ida-----	50	Restricted permeability	0.50	Seepage	0.50
		Slope		Slope	
Monona-----	50	Very limited	1.00	Very limited	1.00
		Slope		Slope	
MsG: Ida-----	50	Restricted permeability	0.50	Seepage	0.50
		Slope		Slope	
Monona-----	50	Very limited	1.00	Very limited	1.00
		Slope		Slope	
On: Onawa-----	100	Restricted permeability	0.50	Seepage	0.50
		Flooding		Flooding	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
				Seepage	1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Pa: Percival-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Pb: Pits-----	100	Not rated		Not rated	
Pc: Platte-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00
PdD2: Ida-----	50	Somewhat limited Restricted permeability Slope	0.50 0.16	Very limited Slope Seepage	1.00 0.50
Ponca-----	50	Somewhat limited Restricted permeability Slope	0.50 0.04	Very limited Slope Seepage	1.00 0.50
PdE2: Ida-----	50	Very limited Slope Restricted permeability	1.00 0.50	Very limited Slope Seepage	1.00 0.50
Ponca-----	50	Somewhat limited Slope Restricted permeability	0.96 0.50	Very limited Slope Seepage	1.00 0.50
Ra: Riverwash-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Rk: Rock Outcrop-----	100	Not rated		Not rated	
Rn: Ida Variant-----	100	Very limited Slope Restricted permeability	1.00 0.50	Very limited Slope Seepage	1.00 0.50
Sd: Inavale-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage	1.00 1.00
Sp: Sarpy-----	100	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Flooding Seepage	1.00 1.00
Ss: Nodaway Variant-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 0.84 0.50	Very limited Flooding Seepage Depth to saturated zone	1.00 0.50 0.17
StE2: Steinauer-----	100	Very limited Restricted permeability Slope	1.00 1.00	Very limited Slope	1.00
Wb: Wabash-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Wc: Wabash-----	100	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Wm: Wann-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Wt: Barney-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
zwa: Water-----	100	Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated	

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Flooding Depth to saturated zone Too clayey	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Too clayey Hard to compact  Depth to saturated zone	1.00 1.00 1.00
Ac: Albaton-----	100	Very limited Flooding Depth to saturated zone Too clayey	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Too clayey Hard to compact  Depth to saturated zone	1.00 1.00 1.00
Af: Alda-----	100	Very limited Flooding Depth to saturated zone Seepage  Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage  Depth to saturated zone	1.00 1.00 0.68
Ag: Alda-----	100	Very limited Flooding Depth to saturated zone Seepage  Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage  Depth to saturated zone	1.00 1.00 0.68
Ca: Carr-----	100	Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Somewhat limited Seepage	0.00
Cc: Cass-----	100	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Somewhat limited Seepage	0.50
Cd: Cass-----	100	Very limited Depth to saturated zone Seepage  Too clayey Flooding	1.00 1.00 0.50 0.40	Very limited Depth to saturated zone Seepage  Flooding	1.00 1.00 0.40	Somewhat limited Too clayey  Depth to saturated zone	0.50 0.09
Ce: Cass-----	100	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Somewhat limited Seepage	0.50
Cg: Colo-----	100	Very limited Flooding  Depth to saturated zone Too clayey	1.00 1.00 0.50	Very limited Flooding  Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Hard to compact  Too clayey	1.00 1.00 0.50
Ck: Colo-----	50	Very limited Flooding  Depth to saturated zone Too clayey	1.00 1.00 0.50	Very limited Flooding  Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Hard to compact  Too clayey	1.00 1.00 0.50
Kennebec-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Not limited	
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.96	Very limited Seepage Slope	1.00 0.96	Very limited Seepage Slope Too Sandy	1.00 0.96 0.50
Ed: Eudora-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
Ga: Gibbon-----	100	Very limited		Very limited		Somewhat limited	



SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Gb: Gibbon-----	100	Flooding	1.00	Flooding	1.00	Depth to saturated zone	0.68
		Depth to saturated zone	1.00	Depth to saturated zone	1.00		
		Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.68
		Depth to saturated zone Seepage	1.00	Depth to saturated zone	1.00		
Gc: Gibbon-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.68
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too clayey	0.50
		Too clayey	0.50				
Gs: Gibbon-----	65	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.68
		Depth to saturated zone Too clayey	0.50	Depth to saturated zone	1.00	Too clayey	0.50
	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Sodium content	1.00
		Depth to saturated zone Sodium content Too clayey	1.00 1.00 0.50	Depth to saturated zone	1.00	Depth to saturated zone Too clayey	0.68 0.50
Gu: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ha: Haynie-----	100	Not limited		Not limited		Not limited	
IdD2: Ida-----	100	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63
IdE: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
IdE2: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Im: Inavale-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
In: Inavale-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
JuB: Judson-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ke: Kennebec-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Not limited	
Le: Lockton-----	100	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage Depth to saturated zone	1.00 1.00 0.68
LS: Luton-----	100	Very limited Depth to saturated zone Too clayey Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Too clayey Hard to compact Depth to saturated zone	1.00 1.00
Lt: Luton-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Too clayey Hard to compact	1.00 1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lu: Luton-----	100	Flooding	0.40			Depth to saturated zone	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too clayey	1.00
		Too clayey	1.00	Flooding	0.40	Hard to compact Depth to saturated zone	1.00
		Flooding	0.40				1.00
MaA: Marshall-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
MaB: Marshall-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
MaC: Marshall-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
MaC2: Marshall-----	100	Not limited		Not limited		Not limited	
MaD: Marshall-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
MeD2: Marshall-----	70	Somewhat limited Too clayey	0.50	Somewhat limited Slope	0.04	Somewhat limited Too clayey	0.50
Ponca-----	30	Slope	0.04	Slope	0.04	Slope	0.04
		Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
MfE: Marshall-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
MfE2: Marshall-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
MoA: Monona-----	100	Not limited		Not limited		Not limited	
MoB: Monona-----	100	Not limited		Not limited		Not limited	
MoC: Monona-----	100	Not limited		Not limited		Not limited	
MoC2: Pohocco-----	100	Not limited		Not limited		Not limited	
MoD: Monona-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
MoD2: Pohocco-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
MoE: Monona-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
MsE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Pohocco-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
		Too clayey	0.50			Too clayey	0.50
MsF: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
MsF2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
MsG: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
On: Onawa-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.68

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Pa: Percival-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	0.00
		Seepage	1.00	Seepage	1.00		
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
Pb: Pits-----	100	Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.68
		Too Sandy	1.00				
		Not rated		Not rated		Not rated	
Pc: Platte-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Too Sandy	1.00	Seepage	1.00	Depth to saturated zone	0.68
		Seepage	1.00				
PdD2: Ida-----	50	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16
Ponca-----	50	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04
PdE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96
Ra: Riverwash-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Gravel content	0.30
Rk: Rock Outcrop-----	100	Not rated		Not rated		Not rated	
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Sd: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00				
Sp: Sarpy-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00				
Ss: Nodaway Variant----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00		
StE2: Steinauer-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Too clayey	0.50			Hard to compact Too clayey	1.00
							0.50
Wb: Wabash-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too clayey	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Hard to compact	1.00
		Too clayey	1.00			Depth to saturated zone	1.00
Wc: Wabash-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too clayey	1.00

SANITARY FACILITIES--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Wm: Wann-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Hard to compact	1.00
		Too clayey	1.00			Depth to saturated zone	1.00
		Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Seepage	0.50
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.44
Wt: Barney-----	100	Seepage	1.00	Seepage	1.00		
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	



AGRICULTURAL WASTE MANAGEMENT  
Douglas and Sarpy Counties, Nebraska

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT  
Douglas and Sarpy Counties, Nebraska

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ab: Albaton-----	100	Very limited Restricted permeability Depth to saturated zone Flooding Runoff limitation	1.00 1.00 0.60 0.40	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 1.00	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.60
Ac: Albaton-----	100	Very limited Restricted permeability Depth to saturated zone Flooding Runoff limitation	1.00 1.00 0.60 0.40	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 1.00	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.60
Af: Alda-----	100	Very limited Filtering capacity Depth to saturated zone Flooding Droughty Sodium content	1.00 0.95 0.60 0.43 0.08	Very limited Filtering capacity Flooding Depth to saturated zone Droughty Sodium content	1.00 1.00 0.95 0.43 0.08	Very limited Filtering capacity Depth to saturated zone Flooding Droughty Sodium content	1.00 0.95 0.60 0.43 0.08
Ag: Alda-----	100	Very limited Filtering capacity Depth to saturated zone Flooding Droughty Sodium content	1.00 0.95 0.60 0.17 0.08	Very limited Filtering capacity Flooding Depth to saturated zone Droughty Sodium content	1.00 1.00 0.95 0.17 0.08	Very limited Filtering capacity Depth to saturated zone Flooding Droughty Sodium content	1.00 0.95 0.60 0.17 0.08
Ca: Carr-----	100	Somewhat limited Flooding Filtering capacity	0.60 0.00	Very limited Flooding Filtering capacity	1.00 0.00	Somewhat limited Flooding Filtering capacity	0.60 0.00
Cc: Cass-----	100	Very limited Filtering capacity Too acid	1.00 0.05	Very limited Filtering capacity Flooding Too acid	1.00 0.40 0.21	Very limited Filtering capacity Too acid	1.00 0.21
Cd: Cass-----	100	Very limited Filtering capacity Depth to saturated zone Restricted permeability Too acid	1.00 0.43 0.41 0.05	Very limited Filtering capacity Depth to saturated zone Flooding Restricted permeability Too acid	1.00 0.43 0.40 0.31 0.21	Very limited Filtering capacity Depth to saturated zone Restricted permeability Too acid	1.00 0.43 0.31 0.21
Ce: Cass-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Filtering capacity	1.00
Cg: Colo-----	100	Very limited Depth to saturated zone Leaching limitation Flooding Restricted permeability	1.00 0.70 0.60 0.41	Very limited Depth to saturated zone Flooding Restricted permeability	1.00 1.00 0.31	Very limited Depth to saturated zone Flooding Restricted permeability	1.00 0.60 0.31
Ck: Colo-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00



AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Kennebec-----	50	Leaching limitation	0.70	Flooding	1.00	Flooding	0.60
		Flooding	0.60	Restricted permeability	0.31	Restricted permeability	0.31
		Restricted permeability	0.41				
		Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
Cm: Urban Land-----	100	Not rated		Not rated		Not rated	
DcE: Dickinson-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Slope	0.96	Slope	0.96	Filtering capacity Too steep for sprinkler application	1.00
							0.97
Ed: Eudora-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
Ga: Gibbon-----	100	Very limited Filtering capacity	1.00	Very limited Flooding	1.00	Very limited Filtering capacity	1.00
		Depth to saturated zone	0.95	Filtering capacity	1.00	Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Too acid	0.91
		Too acid	0.32	Too acid Low adsorption	0.91 0.47	Flooding	0.60
Gb: Gibbon-----	100	Somewhat limited Depth to saturated zone	0.95	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
Gc: Gibbon-----	100	Somewhat limited Depth to saturated zone	0.95	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
		Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
Gs: Gibbon-----	65	Somewhat limited Depth to saturated zone	0.95	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
		Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
Saltine-----	35	Very limited Sodium content	1.00	Very limited Flooding	1.00	Very limited Sodium content	1.00
		Depth to saturated zone	0.95	Sodium content	1.00	Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
		Salinity	0.50	Restricted permeability	0.31	Restricted permeability	0.31
		Restricted permeability	0.41	Salinity	0.00	Salinity	0.00
Gu: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
						Too steep for sprinkler application	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ha: Haynie-----	100	Not limited		Not limited		Not limited	
IdD2: Ida-----	100	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Too steep for surface application Too steep for sprinkler application	1.00  0.77
IdE: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application Too steep for sprinkler application	1.00  1.00
IdE2: Ida-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application Too steep for sprinkler application	1.00  1.00
Im: Inavale-----	100	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.87 0.45	Very limited Filtering capacity Droughty	1.00 0.87	Very limited Filtering capacity Droughty	1.00 0.87
In: Inavale-----	100	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.94 0.45	Very limited Filtering capacity Droughty	1.00 0.94	Very limited Filtering capacity Droughty Too steep for surface application	1.00 0.94 0.31
JuB: Judson-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Too steep for surface application Restricted permeability	0.31 0.31
Ke: Kennebec-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
Le: Lockton-----	100	Very limited Filtering capacity Depth to saturated zone Flooding	1.00 0.95 0.60	Very limited Filtering capacity Flooding Depth to saturated zone	1.00 1.00 0.95	Very limited Filtering capacity Depth to saturated zone Flooding	1.00 0.95 0.60
Ls: Luton-----	100	Very limited Restricted permeability Depth to saturated zone Runoff limitation	1.00 1.00 0.40	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.40	Very limited Restricted permeability Depth to saturated zone	1.00 1.00
Lt: Luton-----	100	Very limited Restricted permeability Depth to saturated zone Runoff limitation	1.00 1.00 0.40	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.40	Very limited Restricted permeability Depth to saturated zone	1.00 1.00
Lu: Luton-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
MaA: Marshall-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Runoff limitation	0.40	Flooding	0.40		
MaB: Marshall-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Restricted permeability	0.31
MaC: Marshall-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Restricted permeability	0.31
MaC2: Marshall-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Too steep for surface application Restricted permeability	0.31
MaD: Marshall-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Too steep for surface application Restricted permeability	0.31
		Slope	0.04	Slope	0.04	Very limited Too steep for surface application Restricted permeability Too steep for sprinkler application	1.00
MeD2: Marshall-----	70	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Restricted permeability Too steep for sprinkler application	0.31
		Slope	0.04	Slope	0.04	Very limited Too steep for surface application Restricted permeability Too steep for sprinkler application	0.22
Ponca-----	30	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Very limited Too steep for surface application Restricted permeability Too steep for sprinkler application	1.00
		Slope	0.04	Slope	0.04	Restricted permeability Too steep for sprinkler application	0.31
MfE: Marshall-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Too steep for surface application Too steep for sprinkler application Restricted permeability	1.00
		Restricted permeability	0.41	Restricted permeability	0.31	Too steep for sprinkler application Restricted permeability	0.97
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Restricted permeability Too steep for sprinkler application Restricted permeability	0.31
		Restricted permeability	0.41	Restricted permeability	0.31	Very limited Too steep for surface application Too steep for sprinkler application Restricted permeability	1.00
MfE2: Marshall-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Too steep for sprinkler application Restricted permeability	0.97
						Very limited Too steep for surface application	0.31
						Too steep for surface application	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ponca-----	50	Restricted permeability	0.41	Restricted permeability	0.31	Too steep for sprinkler application	0.97
		Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Restricted permeability	0.31
		Restricted permeability	0.41	Restricted permeability	0.31	Very limited Too steep for surface application	1.00
MoA: Monona-----	100	Not limited		Not limited		Too steep for sprinkler application	0.97
MoB: Monona-----	100	Not limited		Not limited		Restricted permeability	0.31
MoC: Monona-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
MoC2: Pohocco-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Somewhat limited Too steep for surface application	0.31
MoD: Monona-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Restricted permeability	0.31
MoD2: Pohocco-----	100	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Very limited Too steep for surface application	1.00
		Slope	0.04	Slope	0.04	Too steep for sprinkler application	0.22
MoE: Monona-----	100	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Too steep for surface application	1.00
MsE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Too steep for sprinkler application	0.97
Pohocco-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Too steep for surface application	1.00
		Restricted permeability	0.41	Restricted permeability	0.31	Too steep for sprinkler application	1.00
MsF: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Restricted permeability	0.31
						Very limited Too steep for surface application	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Too steep for sprinkler application	1.00
MsF2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Too steep for sprinkler application	1.00
MsG: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
Monona-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Too steep for sprinkler application	1.00
On: Onawa-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Too steep for surface application	1.00
		Depth to saturated zone	0.95	Flooding	1.00	Too steep for sprinkler application	1.00
		Flooding	0.60	Depth to saturated zone	0.95	Too steep for sprinkler application	1.00
		Runoff limitation	0.40				
Pa: Percival-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Too steep for surface application	1.00
		Filtering capacity	1.00	Flooding	1.00	Too steep for sprinkler application	1.00
		Depth to saturated zone	0.95	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Flooding	0.60	Depth to saturated zone	0.95	Too steep for sprinkler application	1.00
		Droughty	0.26	Droughty	0.26	Too steep for sprinkler application	1.00
Pb: Pits-----	100	Not rated		Not rated		Too steep for sprinkler application	1.00
Pc: Platte-----	100	Somewhat limited Depth to saturated zone	0.95	Very limited Flooding	1.00	Too steep for sprinkler application	1.00
		Droughty	0.78	Depth to saturated zone	0.95	Too steep for sprinkler application	1.00
		Flooding	0.60	Droughty	0.78	Too steep for sprinkler application	1.00
PdD2: Ida-----	50	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Too steep for surface application	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ponca-----	50	Somewhat limited Restricted permeability	0.41	Somewhat limited Restricted permeability	0.31	Too steep for sprinkler application	0.39
						Very limited Too steep for surface application	1.00
		Slope	0.04	Slope	0.04	Restricted permeability	0.31
						Too steep for sprinkler application	0.22
PdE2: Ida-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
Ponca-----	50	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Too steep for sprinkler application	1.00
						Very limited Too steep for surface application	1.00
		Restricted permeability	0.41	Restricted permeability	0.31	Too steep for sprinkler application	0.97
						Restricted permeability	0.31
Ra: Riverwash-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Rk: Rock Outcrop-----	100	Flooding	1.00	Flooding	1.00	Flooding	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45				
Rn: Ida Variant-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
Sd: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Too steep for sprinkler application	1.00
						Very limited Flooding	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.29	Droughty	0.29
Sp: Sarpy-----	100	Very limited Filtering capacity	1.00	Very limited Flooding	1.00	Very limited Filtering capacity	1.00
Ss: Nodaway Variant----	100	Droughty	0.62	Filtering capacity	1.00	Droughty	0.62
		Flooding	0.60	Droughty	0.62	Flooding	0.60
		Leaching limitation	0.45				
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
StE2: Steinauer-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Douglas and Sarpy Counties, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Wb: Wabash-----	100	Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Restricted permeability	0.30	Restricted permeability	0.22	Too steep for sprinkler application	1.00
						Restricted permeability	0.22
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
Wc: Wabash-----	100	Flooding	0.60	Flooding	1.00	Flooding	0.60
		Runoff limitation	0.40	Too acid	0.21	Too acid	0.21
		Too acid	0.05				
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
Wm: Wann-----	100	Flooding	0.60	Flooding	1.00	Flooding	0.60
		Runoff limitation	0.40	Too acid	0.21	Too acid	0.21
		Too acid	0.05				
		Somewhat limited Depth to saturated zone	0.84	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.84
		Flooding	0.60	Depth to saturated zone	0.84	Flooding	0.60
Wt: Barney-----	100	Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	1.00	Flooding	1.00
		Depth to dense layer	1.00	Droughty	0.63	Droughty	0.63
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	





In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Douglas and Sarpy Counties,  
Nebraska

PAGE 2 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Ab: ALBATON SILT LOAM, OVERWASH	ALBATON	Yes	flood plain	2B3	YES	NO	NO
Ac: ALBATON SILTY CLAY	ALBATON	Yes	flood plain	2B3	YES	NO	NO
Af: ALDA FINE SANDY LOAM	ALDA	No	flood plain	---	---	---	---
Ag: ALDA VERY FINE SANDY LOAM	ALDA	No	flood plain	---	---	---	---
Ca: CARR FINE SANDY LOAM	CARR	No	flood plain	---	---	---	---
Cc: CASS FINE SANDY LOAM	CASS	No	flood plain	---	---	---	---
Cd: CASS FINE SANDY LOAM, LOAMY SUBSTRATUM	CASS	No	flood plain	---	---	---	---
Ce: CASS VERY FINE SANDY LOAM	CASS	No	flood plain	---	---	---	---
Cg: COLO SILTY CLAY LOAM	COLO	Yes	flood plain	2B3	YES	NO	NO
Ck: COLO AND KENNEBEC SOILS	COLO	Yes	flood plain	2B3	YES	NO	NO
	KENNEBEC	No	flood plain	---	---	---	---
Cm: CUT AND FILL LAND	URBAN LAND	Unranked	drainageway, hill	---	---	---	---
DcE: DICKINSON SOILS, 11 TO 17 PERCENT SLOPES	DICKINSON	No	bluff	---	---	---	---
Ed: EUDORA SILT LOAM	EUDORA	No	flood plain	---	---	---	---
Ga: GIBBON LOAMY SAND, OVERWASH	GIBBON	No	flood plain	---	---	---	---
Gb: GIBBON SILT LOAM	GIBBON	No	flood plain	---	---	---	---
	PONDED SOILS	Yes	oxbow	2B3,3	YES	NO	YES
Gc: GIBBON SILTY CLAY LOAM	GIBBON	No	flood plain	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
Gs: GIBBON-SLICKSPOTS COMPLEX	GIBBON	No	flood plain	---	---	---	---
	SALTINE WT AT 0-1 FOOT	Yes	flood plain swale	2B3	YES	NO	NO
Gu: GULLIED LAND	IDA VARIANT	No	loess hill	---	---	---	---
Ha: HAYNIE SILT LOAM	HAYNIE	No	flood plain	---	---	---	---
IdD2: IDA SILT LOAM, 7 TO 17 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
IdE: IDA SILT LOAM, 17 TO 30 PERCENT SLOPES	IDA	No	loess hill	---	---	---	---
IdE2: IDA SILT LOAM, 17 TO 30 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
Im: INAVALE LOAMY FINE SAND	INAVALE	No	flood plain	---	---	---	---
	PONDED SOILS	Yes	oxbow	2B3,3	YES	NO	YES
In: INAVALE LOAMY FINE SAND, HUMMOCKY	INAVALE	No	flood plain	---	---	---	---
JuB: JUDSON SILT LOAM, 3 TO 7 PERCENT SLOPES	JUDSON	No	drainageway	---	---	---	---
	WT AT 0-1 FOOT	Yes	flood plain	2B3	YES	NO	NO
Ke: KENNEBEC SILT LOAM, OCCASIONALLY FLOODED	KENNEBEC	No	flood plain	---	---	---	---
	PONDED SOILS	Yes	oxbow	2B3,3	YES	NO	YES

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Douglas and Sarpy Counties,  
Nebraska

PAGE 3 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Le: LEX SOILS, NONCALCAREOUS VARIANT	LOCKTON	No	flood plain	---	---	---	---
	WT AT 0-1 FOOT	Yes	oxbow	2B3	YES	NO	NO
Ls: LUTON SILT LOAM, OVERWASH	LUTON	Yes	flood plain	2B3	YES	NO	NO
Lt: LUTON SILTY CLAY LOAM	LUTON	Yes	flood plain	2B3	YES	NO	NO
Lu: LUTON SILTY CLAY	LUTON	Yes	flood plain	2B3	YES	NO	NO
MaA: MARSHALL SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	MARSHALL	No	loess hill	---	---	---	---
MaB: MARSHALL SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES	MARSHALL	No	loess hill	---	---	---	---
MaC: MARSHALL SILTY CLAY LOAM, 3 TO 7 PERCENT SLOPES	MARSHALL	No	loess hill	---	---	---	---
MaC2: MARSHALL SILTY CLAY LOAM, 3 TO 7 PERCENT SLOPES, ERODED	MARSHALL	No	loess hill	---	---	---	---
MaD: MARSHALL SILTY CLAY LOAM, 7 TO 11 PERCENT SLOPES	MARSHALL	No	loess hill	---	---	---	---
MeD2: MARSHALL-PONCA SILTY CLAY LOAMS, 7 TO 11 PERCENT SLOPES, ERODED	MARSHALL	No	loess hill	---	---	---	---
	PONCA	No	loess hill	---	---	---	---
MfE: MARSHALL AND PONCA SOILS, 11 TO 17 PERCENT SLOPES	MARSHALL	No	loess hill	---	---	---	---
	PONCA	No	loess hill	---	---	---	---
MfE2: MARSHALL AND PONCA SOILS, 11 TO 17 PERCENT SLOPES, ERODED	MARSHALL	No	loess hill	---	---	---	---
	PONCA	No	loess hill	---	---	---	---
MoA: MONONA SILT LOAM, 0 TO 1 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
MoB: MONONA SILT LOAM, 1 TO 3 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
MoC: MONONA SILT LOAM, 3 TO 7 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
MoC2: MONONA SILT LOAM, 3 TO 7 PERCENT SLOPES, ERODED	POHOCCO	No	loess hill	---	---	---	---
MoD: MONONA SILT LOAM, 7 TO 11 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
MoD2: MONONA SILT LOAM, 7 TO 11 PERCENT SLOPES, ERODED	POHOCCO	No	loess hill	---	---	---	---
MoE: MONONA SILT LOAM, 11 TO 17 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
MsE2: MONONA AND IDA SILT LOAMS, 11 TO 17 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
	POHOCCO	No	loess hill	---	---	---	---
MsF: MONONA AND IDA SILT LOAMS, 17 TO 30 PERCENT SLOPES	IDA	No	loess hill	---	---	---	---
	MONONA	No	loess hill	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Douglas and Sarpy Counties,  
Nebraska

PAGE 4 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
MsF2: MONONA AND IDA SILT LOAMS, 17 TO 30 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
MsG: MONONA AND IDA SILT LOAMS, 30 TO 60 PERCENT SLOPES	MONONA	No	loess hill	---	---	---	---
	IDA	No	loess bluff	---	---	---	---
On: ONAWA SILTY CLAY	ONAWA	No	loess hill	---	---	---	---
Pa: PERCIVAL SILTY CLAY	ALBATON	No	flood plain	---	---	---	---
Pb: PITS AND DUMPS	PERCIVAL	Yes	flood plain	2B3	YES	NO	NO
Pc: PLATTE SOILS	PITS	No	flood plain	---	---	---	---
	PLATTE WT AT 0-1 FOOT	Unranked	---	---	---	---	---
		Yes	oxbow	2B3	YES	NO	NO
PdD2: PONCA AND IDA SILT LOAMS, 7 TO 11 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
	PONCA	No	loess hill	---	---	---	---
PdE2: PONCA AND IDA SILT LOAMS, 11 TO 17 PERCENT SLOPES, ERODED	IDA	No	loess hill	---	---	---	---
	PONCA	No	loess hill	---	---	---	---
Ra: RIVERWASH	RIVERWASH	Unranked	flood plain	---	---	---	---
Rk: ROCK LAND	ROCK OUTCROP	Unranked	bluff	---	---	---	---
Rn: ROUGH BROKEN LAND, LOESS	IDA VARIANT	Yes	oxbow	2B3,3	YES	NO	YES
Sd: SANDY ALLUVIAL LAND	INAVAL	No	flood plain	---	---	---	---
	WT AT 0-1 FOOT	Yes	oxbow	2B3	YES	NO	NO
Sp: SARPY FINE SAND	SARPY	No	flood plain	---	---	---	---
Ss: SILTY ALLUVIAL LAND	NODAWAY VARIANT PONDED SOILS	No	flood plain	---	---	---	---
StE2: STEINAUER CLAY LOAM, 11 TO 30 PERCENT SLOPES, ERODED	STEINAUER	Yes	oxbow	2B3,3	YES	NO	YES
Wb: WABASH SILT LOAM	WABASH	No	hill	---	---	---	---
Wc: WABASH SILTY CLAY	WABASH	Yes	flood plain	2B3	YES	NO	NO
Wm: WANN FINE SANDY LOAM	WANN	Yes	flood plain	2B3	YES	NO	NO
	PONDED SOILS	No	flood plain	---	---	---	---
		Yes	oxbow	2B3,3	YES	NO	YES
Wt: WET ALLUVIAL LAND	BARNEY	Yes	oxbow	2B3	YES	NO	NO
zwa: WATER < 40 ACRES	WATER	Yes	flood plain	2B3	YES	NO	NO
zwb: WATER > 40 ACRES	WATER	Unranked	---	---	---	---	---
		Unranked	---	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Douglas and Sarpy Counties,  
Nebraska

PAGE 5 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II.

Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
  - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
  - b. poorly drained or very poorly drained and have either:
    - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in),  
or for other soils
    - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
    - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.



HIGHLY ERODIBLE LANDS REPORT

Survey Area- DOUGLAS AND SARPY COUNTIES, NEBRASKA

Map Symbol	Soil Mapunit Name	HEL Classifications		
		C=10	R=150	wnd wat mu
Ab	ALBATON SILT LOAM, OVERWASH	3	3	3
Ac	ALBATON SILTY CLAY	3	3	3
Af	ALDA FINE SANDY LOAM	3	3	3
Ag	ALDA VERY FINE SANDY LOAM	3	3	3
Ca	CARR FINE SANDY LOAM	3	3	3
Cc	CASS FINE SANDY LOAM	3	3	3
Cd	CASS FINE SANDY LOAM, LOAMY SUBSTRATUM	3	3	3
Ce	CASS VERY FINE SANDY LOAM	3	3	3
Cg	COLO SILTY CLAY LOAM	3	3	3
Ck	COLO AND KENNEBEC SOILS	3	3	3
Cm	CUT AND FILL LAND		2	2
DcE	DICKINSON SOILS, 11 TO 17 PERCENT SLOPES	3	1	1
Ed	EUDORA SILT LOAM	3	3	3
Ga	GIBBON LOAMY SAND, OVERWASH	3	3	3
Gb	GIBBON SILT LOAM	3	3	3
Gc	GIBBON SILTY CLAY LOAM	3	3	3
Gs	GIBBON-SLICKSPOTS COMPLEX	3	3	3
Gu	GULLIED LAND	3	1	1
Ha	HAYNIE SILT LOAM	3	3	3
IdD2	IDA SILT LOAM, 7 TO 17 PERCENT SLOPES, ERODED	3	1	1
IdE	IDA SILT LOAM, 17 TO 30 PERCENT SLOPES	3	1	1
IdE2	IDA SILT LOAM, 17 TO 30 PERCENT SLOPES, ERODED	3	1	1
Im	INAVALE LOAMY FINE SAND	3	3	3
In	INAVALE LOAMY FINE SAND, HUMMOCKY	3	2	2
JuB	JUDSON SILT LOAM, 3 TO 7 PERCENT SLOPES	3	2	2
Ke	KENNEBEC SILT LOAM, OCCASIONALLY FLOODED	3	3	3
Le	LEX SOILS, NONCALCAREOUS VARIANT	3	3	3
Ls	LUTON SILT LOAM, OVERWASH	3	3	3
Lt	LUTON SILTY CLAY LOAM	3	3	3
Lu	LUTON SILTY CLAY	3	3	3
MaA	MARSHALL SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
MaB	MARSHALL SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES	3	3	3
MaC	MARSHALL SILTY CLAY LOAM, 3 TO 7 PERCENT SLOPES	3	2	2
MaC2	MARSHALL SILTY CLAY LOAM, 3 TO 7 PERCENT SLOPES, ERODED	3	2	2
MaD	MARSHALL SILTY CLAY LOAM, 7 TO 11 PERCENT SLOPES	3	1	1
MeD2	MARSHALL-PONCA SILTY CLAY LOAMS, 7 TO 11 PERCENT SLOPES, ERODED	3	1	1
MfE	MARSHALL AND PONCA SOILS, 11 TO 17 PERCENT SLOPES	3	1	1
MfE2	MARSHALL AND PONCA SOILS, 11 TO 17 PERCENT SLOPES, ERODED	3	1	1
MoA	MONONA SILT LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
MoB	MONONA SILT LOAM, 1 TO 3 PERCENT SLOPES	3	3	3
MoC	MONONA SILT LOAM, 3 TO 7 PERCENT SLOPES	3	2	2
MoC2	MONONA SILT LOAM, 3 TO 7 PERCENT SLOPES, ERODED	3	2	2
MoD	MONONA SILT LOAM, 7 TO 11 PERCENT SLOPES	3	1	1
MoD2	MONONA SILT LOAM, 7 TO 11 PERCENT SLOPES, ERODED	3	1	1
MoE	MONONA SILT LOAM, 11 TO 17 PERCENT SLOPES	3	1	1
MsE2	MONONA AND IDA SILT LOAMS, 11 TO 17 PERCENT SLOPES, ERODED	3	1	1
MsF	MONONA AND IDA SILT LOAMS, 17 TO 30 PERCENT SLOPES	3	1	1
MsF2	MONONA AND IDA SILT LOAMS, 17 TO 30 PERCENT SLOPES, ERODED	3	1	1
MsG	MONONA AND IDA SILT LOAMS, 30 TO 60 PERCENT SLOPES	3	1	1
On	ONAWA SILTY CLAY	3	3	3
Pa	PERCIVAL SILTY CLAY	3	3	3
Pb	PITS AND DUMPS	3	2	2
Pc	PLATTE SOILS	3	3	3
PdD2	PONCA AND IDA SILT LOAMS, 7 TO 11 PERCENT SLOPES, ERODED	3	1	1

PdE2	PONCA AND IDA SILT LOAMS, 11 TO 17 PERCENT SLOPES, ERODED	3	1	1
Ra	RIVERWASH		2	2
Rk	ROCK LAND		2	2
Rn	ROUGH BROKEN LAND, LOESS	3	1	1
Sd	SANDY ALLUVIAL LAND	3	3	3
Sp	SARPY FINE SAND	3	3	3
Ss	SILTY ALLUVIAL LAND	3	3	3
StE2	STEINAUER CLAY LOAM, 11 TO 30 PERCENT SLOPES, ERODED	3	1	1
Wb	WABASH SILT LOAM	3	3	3
Wc	WABASH SILTY CLAY	3	3	3
Wm	WANN FINE SANDY LOAM	3	3	3
Wt	WET ALLUVIAL LAND	3	3	3